

Hazardous chemicals in branded textile products on sale in 27 places during 2012

Kevin Brigden, Iryna Labunska, Emily House, David Santillo & Paul Johnston

Greenpeace Research Laboratories Technical Report 06/2012

Contents

Executive Summary	2
1. Introduction	4
2. Materials and methods	5
2.1 Nonylphenol ethoxylates (NPEs)	5
2.2 Carcinogenic amines released under reducing conditions	5
2.3 Phthalates in plastisol printes	5
2.4 Chemical screening	5
3. Results and discussion	6
3.1 Nonylphenol ethoxylates (NPEs)	6
3.2 Carcinogenic amines released under reducing conditions	8
3.3 Phthalates in plastisol prints	9
3.4 Chemical screening	12
4. Conclusions	15
5. References	17
 Table A1: Concentrations of NPEs, carcinogenic amines and phthalates in all articles tested	 22
Table A2: Concentrations of individual phthalates in the 31 articles tested	26
Table A3: Additional substances identified using qualitative chemical screening	27

Executive Summary

Textile manufacturing makes use of a diverse range of process and finishing chemicals, some of which have intrinsic hazardous properties. As a result, finished textile products can contain certain hazardous chemicals used during their manufacture. This may be because of their use as components of the products themselves, or due to residues remaining from their use within the manufacturing processes. Detection of chemical additives and residues in finished textiles can, therefore, provide an indication of chemicals used and potentially released during manufacture.

This report follows on from, and extends, a previous investigation by Greenpeace International that found nonylphenol ethoxylate (NPE) residues to be widespread in a range of sports and recreational textile products. NPEs are used in many countries in the manufacture of textiles, in particular as surfactants and detergents, although such uses have effectively been banned within the EU, with similar restrictions in place in the US and Canada. Once released, NPEs break down to form nonylphenol isomers. These are persistent, bioaccumulative and toxic chemicals.

The current study extended the range of chemicals investigated and broadened the range of items analysed for these chemicals. In all, 141 high street fashion textile products were analysed. These included items sold by 20 different major clothing brands, purchased in 27 countries around the world. The range included examples manufactured in at least 18 different countries, although the countries of manufacture of 25 of the articles were unknown. The investigation included products designed for men, women and children, and covered a variety of articles including t-shirts, jeans/trousers, dresses and underwear, as well as various other types of clothing.

As in the previous investigation, the concentration of NPEs was determined in all items. In addition, all items that included dyed fabric (134 samples) were tested for carcinogenic amines known to be released from azo dyes under certain conditions. In addition, toxic phthalate esters (commonly referred to as phthalates) were quantified in 31 items that featured medium or large plastisol printed areas. Moreover, just under half of the products (63 samples) were investigated using a broader qualitative chemical screening to identify the presence, as far as possible, of any residues of other chemicals present extractable in each product. Information on the hazardous properties of the chemicals quantified in this study, together with those identified through chemical screening, is provided within the report.

Key findings of this study include:-

- **NPEs** were the most commonly detected substances, identified in 89 of the 141 articles (63% of the total). NPE concentrations ranged from just above 1 mg/kg up to 45 000 mg/kg. One fifth (20%) of the samples contained NPEs at concentrations above 100 mg/kg. Of this 20%, 12 samples (9% of articles tested) showed concentrations over 1 000 mg/kg (0.1% by mass),
- NPEs were detected in one or more products from all of the brands included in this study. NPEs were also found in at least one product from 13 of the 18 identifiable countries of manufacture, and in products sold in 25 out of the 27 countries. Overall, the NPE results were consistent with those from the previous study carried out by Greenpeace International in which 78 items of sports and recreational clothing and shoes were analysed.
- **Phthalates** were detected in all 31 samples of fabric bearing a plastisol print. For four of these samples, phthalates were present at high or very high concentrations, indicating their use as plasticisers within the plastisol formulations. Three samples contained total phthalate levels of between 20.0% and 37.6% by weight (200 000 to 376 000 mg/kg), and a fourth contained 0.52% by weight (5 200 mg/kg). The origins of the phthalates detected in the other 27 samples with lower total phthalate levels (4 to 138 mg/kg) are not certain, and sources other than use during manufacture may contribute. Nonetheless, the presence of phthalates within a product at any level is of concern whatever the source,
- The individual phthalates present in the four samples with high or very high levels were DEHP (di-2-ethylhexyl phthalate), BBP (butyl benzyl phthalate) or DINP (diisononyl phthalate),
- A **carcinogenic amine**, *o*-dianisidine, was released from two items under the test conditions, at 7 mg/kg and 9 mg/kg respectively. Any release of such a carcinogenic compound is of concern, though the levels of *o*-dianisidine were below regulatory limits set in certain countries (including the EU and China) designed to prohibit the sale of textile products containing azocolourants that can release carcinogenic amines under the test conditions,
- Numerous other industrial chemicals were identified in various individual items, including many that are hazardous or potentially hazardous, some of which are classified as toxic or very toxic to aquatic life.

Overall, a variety of hazardous chemicals were detected within the broad range of high street fashion textile products analysed. These covered a diverse range of brands and countries of manufacture. These results indicate the ongoing – and in some cases widespread – use of hazardous chemicals in the manufacture of textile products openly marketed to consumers. Specifically, the presence of NPEs within a wide range of items indicates that NPE use is widespread within the international textile industry, including within the supply chains used by many major international clothing brands. The presence of phthalates at high levels in some items shows that the presence of hazardous chemicals within some items is due to their intentional incorporation into certain textile products, and is not only due to residues left over from use in manufacturing processes.

In the case of the NPEs, which are readily water-soluble, it should be noted that the concentration in a product cannot be considered indicative of the amount of NPEs used during manufacture. Residue levels will inevitably depend on a range of other factors, including the number of wash cycles employed during product finishing and the efficiency with which these chemicals are removed. For the same reason, the absence of NPE residues from any particular product cannot be taken to confirm absence of use of NPEs at some stage during the manufacturing processes.

The presence of chemicals in textile products indicates both that they are used during manufacture, and that, therefore, there is a potential for their release from manufacturing facilities. It also highlights the potential for releases from the products themselves after they have been sold. An earlier investigation by Greenpeace International has shown that NPEs within textiles are readily washed out when they are laundered. Most, if not all, the other hazardous chemicals identified in this study are also likely to be washed out to some degree when the product is laundered. In addition, phthalates in plastisol formulations are not tightly bound to the plastic, but are present as mobile components within the matrix, and will, therefore, be released from the product over time.

Each individual garment may contain only relatively small quantities of the chemicals identified. Nonetheless, given the overall volumes of textile production and of retail sales, the aggregated quantities of these hazardous chemicals present in textile products could be significant, with total releases representing a significant diffuse source of chemicals in the country of sale.

The range of hazardous or potentially hazardous chemicals identified in the broad screening analysis of 63 items indicates that investigations that focus only on a narrow range of target substances will inevitably underestimate the complexity of chemical residues in finished textiles and, therefore, the overall potential for chemical releases during manufacture and subsequent laundering.

Overall, this study has provided a greater understanding of the presence and, in some cases, concentrations of a broad range of chemicals within high street fashion textile products across a diverse range of brands, and of countries of manufacture and sale. The number of articles investigated in this study is inevitably small compared to the vast number of products manufactured and sold per country or per brand. It is, therefore, not possible to draw more general conclusions about the levels or presence of such chemicals that may be expected for all such products. Rather, this study provides a snapshot of what appears to be a generic problem that is not restricted to any particular country, product type or brand. It is one that deserves further investigation. In particular, the current regulatory frameworks need scrutiny and development to improve their effectiveness in regulating this sector of industrial manufacture and marketing. In addition, suppliers and retailers need to develop robust procurement policies designed to force the elimination of such chemicals from manufacturing processes, and hence, in the finished products.

1. Introduction

Finished textile products can contain certain hazardous chemicals used during their manufacture, either because of their use as components of materials incorporated within the product or due to residues remaining from the use within processes employed during manufacture. In either case, textile products can subsequently act as sources of these chemicals to the environment, especially to water during washing. While each individual garment may release only a relatively small quantity of hazardous chemicals through its lifetime, given the overall volumes of textile production and retail, release from textiles as an overall product sector could represent a significant diffuse source of chemicals to water in the country of sale when considered on a national or regional basis. The presence of chemical additives and residues in finished textiles also gives an indication of the nature of chemical use and release during processes by which they are manufactured.

This report follows on from, and extends, research recently published by Greenpeace International that identified the presence of one group of hazardous chemicals, the hormone-disrupting nonylphenol ethoxylates (NPEs), in a range of textile products consisting primarily of sports and recreational clothing and shoes (Brigden et al. 2012, Greenpeace 2011b).

This current study investigated the presence of hazardous chemicals in a broader range of textile products, consisting of 141 textile products across many countries of manufacture and sale, and for a wide range of major clothing brands. The majority of products were tested quantitatively for NPEs, as well as for carcinogenic amines released from azo dyes within dyed fabric, and phthalate esters (commonly referred to as phthalates) in fabrics bearing a plastisol print¹. In addition, just under half of the products (63 samples) from a selection of the brands included in this study were investigated through a broader qualitative chemical screening to identify the presence, as far as possible, of any other hazardous chemicals present within the products, in order to provide an indication of the extent and nature of other chemical residues within textile products of this type.

As mentioned above, the presence of NPEs has been previously reported in certain textile products (Brigden et al. 2012, Greenpeace 2011b, and references within these reports). Similarly, the presence in certain textile products has also been previously reported for carcinogenic amines released under reducing conditions from azo dyes (JRC 2008, Laursen et al. 2003) and phthalates in plastisol printed fabric (RAPEX 2012, Greenpeace 2004). The use of a chemical screening approach by Laursen et al. (2003) also identified the presence of a wider range of chemicals within a small number of products).

More information on the chemicals investigated in the current study is provided in Boxes A-D.

The 141 products included in this study included items sold by 20 different major clothing brands, and were purchased across 27 countries around the world. According to their labels, the 141 products included examples that were manufactured in at least 18 different countries, with 25 articles being of unknown manufacturing origin. The sample set included products designed for men, women and children, and covered a variety of articles including t-shirts, jeans/trousers, dresses and underwear, as well as a variety of other types of clothing.

¹ **Plastisol:** A suspension of plastic particles, commonly PVC or EVA, in a plasticiser. Used as ink for screen-printing images and logos onto textiles.

2. Materials and methods

The 141 products were purchased in April 2012 at the flagship stores of the clothing brands, or other stores authorised to sell the branded products. While still in the store, purchased products were immediately sealed in individual identical clean polyethylene bags. Sealed bags containing the products were sent to the Greenpeace Research Laboratories at the University of Exeter in the UK, from where they were dispatched to independent accredited laboratories for analysis of NPEs, phthalates and azo dyes-related amines. For just under half of the products, a broader qualitative screening for other chemicals present within the fabric was carried out at the Greenpeace Research Laboratories. Details of the individual articles are provided in Appendix 1.

2.1 Nonylphenol ethoxylates (NPEs)

The concentrations of NPEs were quantified in all 141 articles of clothing. For the majority of articles (110 of 141), a section of fabric that did not bear a plastisol print was tested. For the remaining 31 products, the concentration of NPEs was quantified in a section of fabric bearing a plastisol print of an image, logo or text. These 31 articles were also investigated for the presence of phthalates within the printed fabric (see Section 2.3).

Following isolation of a section of fabric from each article, the sample was extracted with an acetonitrile-water mixture in the ratio 70:30 and then analysed with reversed-phase HPLC liquid chromatography along with Applied Biosystems' API 4000 tandem mass spectrometry (LC-MS/MS). The quantification was carried out for each of 17 individual NPEs, consisting of those with between 4 and 20 ethoxylate groups. The quantitative results presented below are the sum of the concentrations of the individual NPEs with 4-20 ethoxylate groups. Quality assurance and quality control checks were employed, including the analysis of seven samples in duplicate and the analysis of blank samples and two intra-laboratory textile reference samples with every batch, of between 11 to 52 samples.

2.2 Carcinogenic amines released under reducing conditions

With the exception of 7 articles for which the fabric was white in colour, all articles that were investigated for the presence of NPEs were also investigated for the concentrations of carcinogenic amines released under certain reducing conditions, related to the presence of certain azo dyes (see Box B for detail). The 134 articles were tested in accordance with method EN 14362 related to the relevant EU regulations (EU 2002), which involved the determination of certain aromatic amines derived from azo colorants following cleavage of the azo group under reducing conditions, either directly or following extraction from the fabric, depending on the type of fabric in each sample. The analysis included the quantification of two additional amines that are listed under the equivalent Chinese regulation (SAPRC 2012).

2.3 Phthalates in plastisol prints

All articles bearing a medium or large-sized plastisol print of an image, logo or text were investigated for the concentrations of a range of phthalates within the printed fabric. These 31 articles, the same ones for which NPEs were quantified within fabric bearing a plastisol print, were quantified for the concentrations of the following phthalates in a section of printed fabric; dimethyl phthalate (DMP), diethyl phthalate (DEP), di-n-butyl phthalate (DnBP), diisobutyl phthalate (DIBP), butylbenzyl phthalate (BBP), di-2-ethylhexyl phthalate (DEHP), di-n-octyl phthalate (DnOP), diisononyl phthalate (DINP) and diisodecyl phthalate (DIDP).

A portion of each sample bearing a plastisol print was extracted with ethyl acetate:cyclohexane (1:1) using deuterated (D8)-naphthalene as a quality control standard to check extraction efficiency. Blanks were also extracted as part of the quality control checks employed. The concentrations of phthalates in the extracts were subsequently analysed by gas chromatography/mass spectrometry (GC/MS) using a LECO time of flight (TOF) instrument with a programmed temperature vaporizing (PTV) injector and a DB5ms column, using deuterated (D10)-pyrene as an internal standard.

2.4 Chemical screening

For each of the 63 samples, a representative portion was extracted using pentane:acetone (3:1) with a Dionex 350 accelerated solvent extractor (ASE), with deuterated (d8)-naphthalene as an internal standard to check extraction efficiency. Concentrated extracts were cleaned through a Florisil column, eluting with pentane:toluene (95:5). Cleaned extracts were concentrated to 0.5 mL and brominated naphthalene added as an internal standard for the analytical procedure.

The final extract was analysed by gas chromatography/mass spectrometry (GC/MS) using an Agilent 6890 Series II GC with a Restek Rtx-XLB column, linked to an Agilent 5973 inert mass selective detector (MSD) operated in electron impact (EI) mode. Identification of compounds was carried out by matching spectra against both the Wiley 7N and Pesticides libraries, combined with expert judgment to confirm identification. In addition to the use of internal standards, a range of blank samples were extracted and/or analysed with each batch of five samples. Additional details on the chemical screening method are provided in Appendix 4.

3. Results and Discussion

The results for the various substance groups are presented in the following sections. Details of all results for the individual articles are provided in Appendix 1, along with a breakdown of the concentrations of all individual phthalates in the relevant articles in Appendix 2, and a listing of all substances identified in individual samples that were investigated using the chemical screening method in Appendix 3.

3.1 Nonylphenol ethoxylates (NPEs)

Of the 141 articles in which NPEs were quantified, 89 articles (63% of the total) tested positive for the presence of NPEs at concentrations above the limit of detection (1 mg/kg), at levels ranging from just above 1 mg/kg up to 45 000 mg/kg. The highest concentration (45 000 mg/kg) was detected in a sample of fabric bearing a plastisol print from a C&A branded t-shirt manufactured and sold in Mexico (our sample code: TX12041). While some other samples bearing a plastisol print also contained relatively high concentrations of NPEs, this was not always the case, with NPEs not being detected above 1 mg/kg in just under half (14 of 31) of the plastisol printed samples.

One fifth (20%) of the samples contained NPEs at concentrations above 100 mg/kg and, of these, 12 samples (9% of articles tested) had concentrations over 1 000 mg/kg (0.1% by mass). Of the remainder, just over two fifths (43%) of all samples tested contained detectable levels of NPEs at concentrations below 100 mg/kg. A summary of the number of samples containing NPEs within various ranges of concentration is given in Table 1.

NPE concentration range (mg/kg)	Number of samples of 141 analysed (% of samples)
<1	52 samples (37%)
1 – 10	27 samples (19%)
>10 – 100	34 samples (24%)
>100 – 1 000	16 samples (11%)
>1 000	12 samples (9%)

Table 1. The number of samples (of the 141 articles tested) within various NPE concentration ranges.

NPEs were detected in one or more product from all of the brands included in this study. Furthermore, NPEs were detected in one or more product from 13 of the 18 countries of manufacture, and in products sold in 25 out of the 27 countries. For the countries of manufacture for which NPEs were not detected in any of the articles tested, only a relatively small number of articles per country were included in the study; Cambodia (2 articles), Jordan (1 article), Romania (1 article), Spain (1 article), and Tunisia (3 articles). A summary of the results is presented in Tables 2 to 4, which include a breakdown of the results by brand, by place of sale, and by country of manufacture, respectively. These tables include the median values for all samples in each brand, place of sale or country of manufacture, as well as the median value for only those samples in which NPEs were detected in each case.

Brand	Number of samples	Number tested positive	NPE concentration range, when detected (mg/kg)	Median of detected levels (mg/kg)	Median of all levels (mg/kg)
Armani	9	5	1.2 – 43	8.1	1.2
Benetton	9	3	6.3 – 95	11	<1
Blazek	4	2	47 – 330	190	24
C&A	6	5	1.7 – 45 000	63	35
Calvin Klein	8	7	5.6 – 4 000	20	17
Diesel	9	3	6.6 – 710	16	<1
Esprit	9	6	1.1 – 770	47	17
Gap	9	7	1.3 – 920	43	8.6
H&M	6	2	1.6 – 8.7	5.2	<1
Jack & Jones	5	3	4.6 – 2 100	17	4.6
Levi's	11	7	5.7 – 4 100	80	9.7
Mango	10	6	1.3 – 9 800	690	4.3
Marks & Spencer	6	4	84 – 2 090	590	319
Metersbonwe	4	3	140 – 2 100	1 500	795
Only	4	4	5.5 – 730	35	35
Tommy Hilfiger	9	6	3.9 – 500	24	8.6
Vancl	4	4	7.6 – 150	76	76
Vero Moda	5	4	6.3 – 130	38	31
Victoria's Secret	4	2	7.0 – 10	8.5	3.8
Zara	10	6	9.6 – 2 600	27	14

Table 2. The number of samples in which NPEs were identified by product brand, with the NPE concentration ranges, the median NPE concentrations in all samples, and the median values for those samples in which NPEs were detected, for each brand.

Place of sale	Number of samples	Number tested positive	NPE concentration range, when detected (mg/kg)	Median of detected levels (mg/kg)	Median of all levels (mg/kg)
Austria	3	2	7.2 – 8.6	7.9	7.2
Belgium	5	2	27 – 80	54	<1
Canada	4	3	8.6 – 4 000	10	9.3
China (mainland)	16	13	5.5 – 2 600	140	32
Czech Republic	6	2	47 – 330	190	<1
Denmark	6	3	17 – 730	45	8.8
Finland	2	0	-	-	<1
France	5	4	6.3 – 110	36	8.1
Germany	4	4	9.0 – 770	360	358
Hong Kong	2	0	-	-	<1
Hungary	4	2	1.7 – 29	15	1.1
Indonesia	5	4	3.8 – 550	36	5.6
Israel	4	3	1.3 – 43	8.7	<1
Italy	6	3	3.9 – 95	11	2.2
Lebanon	5	5	1.6 – 130	15	15
Mexico	6	5	56 – 45 000	920	713
Netherlands	5	0	-	-	<1
Norway	1	1	38	38	38
Philippines	4	3	20 – 620	600	311
Russia	6	3	6.6 – 17	17	3.6
South Africa	6	4	1.3 – 700	9.9	3.5
Spain	5	4	8.7 – 9 800	28	25
Sweden	2	1	43	43	22
Switzerland	6	3	1.1 – 64	6.9	<1
Taiwan	2	2	9.7 – 79	44	44
Thailand	6	3	32 – 2 090	460	16
Turkey	4	3	18 – 1 500	84	51
UK	5	2	4.8 – 1 400	690	<1
USA	6	5	1.2 – 4 100	26	17

Table 3. The number of samples in which NPEs were identified by the place of sale, with the NPE concentration range, the median NPE concentrations in all samples, and the median values for those samples in which NPEs were detected, for each place.

Place of manufacture	Number of samples	Number tested positive	NPE concentration range, when detected (mg/kg)	Median of detected levels (mg/kg)	Median of all levels (mg/kg)
Bangladesh	12	5	1.3 – 1 500	8.6	<1
Cambodia	2	0	-	-	<1
China (mainland)	34	21	1.3 – 2 600	80	9.9
Egypt	2	2	9.0 – 11	10	10
India	9	6	6.3 – 4 000	27	8.7
Indonesia	8	6	1.2 – 620	12	8.4
Jordan	1	0	-	-	<1
Mexico	6	6	56 – 45 000	2 000	2 000
Morocco	3	2	7.2 – 710	360	7.2
Pakistan	3	3	3.8 – 29	19	19
Philippines	1	1	26	26	26
Romania	1	0	-	-	<1
Spain	1	0	-	-	<1
Sri Lanka	2	2	3.9 – 7.0	5.5	5.5
Thailand	3	1	14	14	<1
Tunisia	3	0	-	-	<1
Turkey	15	7	17 – 9 800	38	<1
Vietnam	12	8	4.8 – 700	34	5.7
Unknown	23	19	1.1 – 2 100	31	18

Table 4. The number of samples in which NPEs were identified by place of manufacture, with the NPE concentration range, the median NPE concentrations in all samples, and the median values for those samples in which NPEs were detected, for each place.

Samples with higher NPE concentrations (above 100 mg/kg) were distributed across a range of places of sale, places of manufacture and product brands. However, it is not possible to draw any general conclusions on the variability of NPE concentrations in articles between the place of sale, place of manufacture, or product brand given the relatively small number of products included in this study, compared to the vast number of products manufactured and sold per place or per brand.

Box A. Nonylphenol ethoxylates (NPEs)

Nonylphenol ethoxylates (NPEs) are a group of chemicals used as surfactants, emulsifiers, dispersants and wetting agents in a variety of applications, including the manufacture of textiles. The use of NPEs during the manufacture of textiles can leave residues of NPEs within the final product which are readily released when the items are washed as part of their normal use (Brigden et al. 2012, Greenpeace 2012).

Where NPEs are released from manufacture facilities or through the laundering of textile products, either directly into surface waters or via wastewater treatment facilities, they can break down to form nonylphenol, a closely-related group of persistent, bioaccumulative and toxic chemicals (OSPAR 2004, Jobling et al. 1996). The manufacture, use and release of NPEs and nonylphenol are regulated in some regions (OSPAR 1998, EU 2003), though uses continue elsewhere. Additional information on nonylphenol and NPEs, including on their hazardous properties, is available in a previous related report (Box C, Greenpeace 2011a).

3.2 Carcinogenic amines released under reducing conditions

Of the 134 articles tested, the presence of one or more carcinogenic amines released under the test conditions was detected in two articles, both manufactured in Pakistan. For all other articles, all carcinogenic amines included in the analysis were not detected above the method detection limit (<5 mg/kg).

For the two articles in which release of an amine was detected, the same compound was detected for both; 3,3'-dimethoxybenzidine (also known as *o*-dianisidine). The other carcinogenic amines included in the analysis were not detected for these two articles (<5 mg/kg). Details of the results for these two articles are given in Table 5 below, with full results for all articles tested provided in Appendix 1. Additional information on carcinogenic amines that can be released from certain azo dyes under reducing conditions, including the specific amine detected in this study (*o*-dianisidine), is provided in Box B.

Sample code	Brand	Sold in	Made in	Type	Colour	3,3'-dimethoxy benzidine	Other amines
TX12128	Zara	Lebanon	Pakistan	Jeans (child)	Dark blue	7 mg/kg	<5 mg/kg
TX12130	Zara	Hungary	Pakistan	Jeans (child)	Dark blue	9 mg/kg	<5 mg/kg

Table 5. Details of articles in which amines were detected under the test conditions.

Although the article sold in Hungary (TX12130) is subject to the EU Directive that regulates the sale of products containing azo colourants that can release carcinogenic amines under specific test conditions (EU 2002), the concentration of *o*-dianisidine recorded was below the regulatory limit set within the EU (30 mg/kg; EU 2002) and, while any detectable presence of such a carcinogenic compound is of concern, would not, therefore, make the sale of the product illegal in the EU. As far as we are aware, there is no relevant legislation covering the presence of these compounds within textiles sold in Lebanon. The concentrations of *o*-dianisidine in this product and in the other in which it was detected (TX12130, on sale in the Lebanon), also fell below the stricter limits set for products sold within China (20 mg/kg, SAPRC 2012). It is not known whether equivalent regulations exist within Pakistan, where both products were manufactured, or in Lebanon, where the second of the two products was on sale.

Box B. Carcinogenic amines released by certain azo dyes

Azo dyes can undergo reduction to release aromatic amines. This reductive release can take place under a number of conditions, including within the body; reduction can occur in many different types of cells, including within intestinal and skin bacteria (Golka et al. 2004, Rafi et al. 1997, IARC 2008). Some, though not all, aromatic amines that can be released from azo dyes have been shown to be carcinogenic (IARC 1998, 1987).

Azo dyes are manufactured using the same amines that can be later released through reduction, and it is therefore possible for commercial azo dye formulation to contain residues of amines used in their manufacture. Furthermore, certain carcinogenic amines have been detected as residues in other amines that are used for azo dye manufacture, providing an additional route for contamination of commercial azo dye formulations with carcinogenic amines (IARC 2008). These sources could contribute to the presence of carcinogenic amines at trace levels within textile products.

Legislation exists in certain countries, including EU member states and China, which prohibits the sale of products containing dyes that can degrade under specific test conditions to form carcinogenic amines at concentration above set limits, for textile articles which may come into direct and contact with human skin. The EU regulation lists 22 compounds, with a limit of 30 mg/kg (EU 2002). The regulation in China sets a limit of 20 mg/kg and lists the same compounds as the EU regulation, as well as two additional compounds (SAPRC 2012).

Animal studies have shown that 3,3'-dimethoxybenzidine (also known as *o*-dianisidine), together with certain other benzidine congeners, can have a carcinogenic effect, increasing tumour incidence in many organs (Haley 1975, Morgan *et al.* 1994, IARC 2008). There is clear evidence that exposure to benzidine-based dyes has caused bladder cancer in humans. However, the carcinogenicity of *o*-dianisidine alone has not been conclusively demonstrated in humans through epidemiological studies, partly due to the manufacture and use *o*-dianisidine having occurred together with other amines that are known human carcinogens, affecting the ability to demonstrate *o*-dianisidine as a causal compound (DHHS 2011). The International Agency for Research on Cancer (IARC) has classified *o*-dianisidine as possibly carcinogenic to humans (class 2B) (IARC 1998), and similarly the US Department of Health and Human Service lists *o*-dianisidine and dyes that are metabolised to *o*-dianisidine as reasonably anticipated to be human carcinogens (DHHS 2011).

3.3 Phthalates in plastisol prints

Of the 31 articles in which a range of phthalates were quantified in plastisol printed fabric, four articles contained one or more individual phthalate at a concentration above 1 000 mg/kg (0.1% by mass, see Table 6 for details). Of these, three samples contained very high levels, with concentrations of individual phthalates between 200 000 mg/kg (20% by weight) and 320 000 mg/kg (32% by weight). The fourth sample (TX12119) had a lower but still relevant level, 5 200 mg/kg (0.52% by weight). For all other articles, individual phthalates were present at concentrations of 120 mg/kg or less, with a method detection limit of 3 mg/kg. Full details of the individual phthalate concentrations for all 31 articles are presented in Appendices 1 and 2.

For the four samples that contained phthalates at levels above 0.1% by mass, the predominant phthalates identified were di-2-ethylhexyl phthalate (DEHP) and diisononyl phthalate (DINP), with one sample (TX12008) also containing butyl benzyl phthalate (BBP) as a substantial component. The presence of these phthalates at the concentrations found indicate their use as plasticisers within the plastisol formulations used to manufacture these products.

In contrast, the concentrations recorded in the remaining 27 samples would be too low to have any significant plasticising function on their own. For these articles, the identified phthalates may be present due to contamination of another substance used within the plastisol formulation (including other plasticisers, inks or pigments), or through other uses of the phthalates within the facilities that manufactured the products. In addition, it cannot be excluded that the presence of phthalates at these levels could have arisen from sources unrelated to chemical use at the facilities that manufactured the products, such as through contact with phthalate-bearing materials subsequent to manufacture, up until the point at which the products were purchased and separately sealed for analysis. Nonetheless, the presence of phthalates within a product at any level is of concern, whatever the source.

Sample code	TX12008	TX12110	TX12115	TX12119
Brand	Armani	Tommy Hilfiger	Tommy Hilfiger	Victoria's Secret
Sold in	Italy	USA	Austria	USA
Made in	Turkey	Philippines	Bangladesh	Sri Lanka
Type	T-shirt	T-shirt	T-shirt	Underwear
Phthalate	mg/kg (% by mass)			
Butyl benzyl phthalate (BBP)	23 000 (2.3%)	-	-	-
Di-2-ethylhexyl phthalate (DEHP)	200 000 (20%)	-	56 000 (5.6%)	5 200 (0.52%)
Diisononyl phthalate (DINP)	-	200 000 (20%)	320 000 (32%)	-
NPEs (for comparison)	<1	41	8.6	7.0

Table 6. Concentrations of individual phthalates (mg/kg) where present in articles above 1 000 mg/kg. Also including are the concentrations of NPEs (mg/kg) for comparison.

The phthalates present at high levels in the four articles listed in Table 6 are toxic compounds. DEHP and BBP are known to exert reproductive toxicity, with DEHP being toxic to the developing reproductive system in mammals (Ema & Miyawaki 2002, Mylchreest et al. 2002, Aso et al. 2005). DEHP and BBP, as well as DNBP and DIBP, have recently been listed as Substances of Very High Concern under the European REACH Regulation (ECHA 2011). DINP, though not among the phthalates of greatest regulatory concern, nonetheless does exhibit toxicity (primarily to the liver and kidney) at high doses. In addition, some hormone-disrupting (anti-androgenic) effects on reproductive development in rats have recently been reported for DINP (Boberg et al. 2011). Further background information on phthalate esters is provided in Box C.

Furthermore, phthalates in plastisol formulations are not tightly bound to the plastic, but are present as mobile components within the matrix, and can therefore be released from the product over time (DoE 1991, Cadogan et al. 1993, Jenke et al. 2006, Fierens et al. 2012, Fasano et al. 2012, Latorre et al. 2012).

Box C. Phthalate esters (Phthalates)

Phthalates (or, more accurately, phthalate diesters) are non-halogenated chemicals with a diversity of uses, dominated by use as plasticisers (or softeners) in plastics, especially PVC (e.g., in cables and other flexible components). Other applications included uses as components of inks, adhesives, sealants, surface coatings and personal care products. Some phthalates are discrete chemicals, such as the well known di-2-ethylhexyl phthalate (DEHP), while others are complex mixtures of isomers, such as diisononyl phthalate (DINP) and diisodecyl phthalate (DIDP). All uses of phthalates, especially the major use as PVC plasticisers, result in large-scale losses to the environment (both indoors and outdoors) during the lifetime of products, and again following disposal, principally because phthalates are not chemically bound but only physically associated to the polymer chains. Phthalates have been found to leach from food packaging materials and contaminate corresponding food products (Fierens et al. 2012, Fasano et al. 2012); from tubing material used for drug products manufacturing (Jenke et al. 2006) and from PVC blood bags, which primarily contained DEHP (Ferri et al. 2012). Moreover, it has been shown that bacteria, which may grow on PVC plastics in wet conditions (e.g., shower curtains), may enhance DEHP leaching from plastic (Latorre et al. 2012). Thus, phthalates are widely found in the indoor environment, including in air and dust (Langer et al. 2010, Otake et al. 2001, Butte & Heinzow 2002, Fromme et al. 2004) at concentrations that commonly reflect the prevalence of plastics and certain textiles within the rooms sampled (Abb et al. 2009). Once plastic products are disposed to municipal landfills, phthalates, particularly DIBP and DnBP, may continue to leach, finally reaching groundwater (Liu et al. 2010).

Phthalates are commonly found in human tissues, including in blood, breast milk and, as metabolites, in urine (Colon et al. 2000, Blount et al. 2000, Silva et al. 2004, Guerranti et al. 2012), with reports of significantly higher levels of intake in children (Koch et al. 2006). In humans and other animals, they are relatively rapidly metabolised to their monoester forms, but these are frequently more toxic than the parent compound (Dalgaard et al. 2001).

Substantial concerns exist with regard to the toxicity of phthalates to wildlife and humans. For example, DEHP, one of the most widely used to date, is known to be toxic to reproductive development in mammals, capable (in its monoester form MEHP) of interfering with development of the testes in early life, thought to be mediated through impacts on testosterone synthesis (Howdeshell et al. 2008, Lin et al. 2008). Even at low doses, exposure to mixtures of phthalates can result in cumulative effects on testicular development in rats (Martino-Andrade et al. 2008). In addition, adverse impacts on female reproductive success in adult rats and on development of the young have been reported following exposure to this chemical (Lovekamp-Swan & Davis 2003, Grande et al. 2006, Gray et al. 2006).

A more recent study (Abdul-Ghani et al. 2012) has shown that both DEHP and DBP can induce gross malformations, damage to DNA and changes in behavioural development when administered to developing chick embryos. The review of Caldwell (2012) highlights recently discovered impacts of DEHP including chromosomal damage, increased cancer progression and changes in gene expression at increasingly lower concentrations. Both DEHP and DBP are classified as “toxic to reproduction” within Europe.

Butylbenzyl phthalate (BBP) and dibutyl phthalate (DBP) have also been reported to exert reproductive toxicity (Ema & Miyawaki 2002, Mylchreest et al. 2002, Aso et al. 2005). Other research has revealed a correlation between phthalate exposure during pregnancy and decreased ano-genital index (distance from the anus to the genitals) in male children (Swan et al. 2005). Decreased AGI correlated with concentrations of four phthalate metabolites, namely mono-ethyl phthalate (MEP), mono-n-butyl phthalate (MBP), monobenzyl phthalate (MBzP), and monoisobutyl phthalate (MIBP). It was also found that DBP can be taken up by crop plants as well impacting on physiology and morphology of some crops during growth (Liao 2006). Other commonly used phthalates, including the isomeric forms DINP and DIDP, are of concern because of observed effects on the liver and kidney, albeit at higher doses. DINP has also been found (Boberg et al. 2011) to exhibit anti-androgenic effects on reproductive development of Wistar rats, though less prominent than DEHP, DBP and BBP, however, further safety evaluation of DINP should be undertaken.

At present, there are relatively few controls on the marketing and use of phthalates, despite their toxicity, the volumes used and their propensity to leach out of products throughout their lifetime. Of the controls which do exist, however, probably the best known is the EU-wide ban on the use of six phthalates in children’s toys and childcare articles, first agreed as an emergency measure in 1999 and finally made permanent in 2005 (EC 2005). While this addresses one important exposure route, exposures through other consumer products have so far largely escaped regulation. Within Europe, three phthalates (DBP, BBP and DEHP) were proposed to be included into the first shortlist of seven substances for which detailed justification and authorisation will be required for any proposed continued uses (ECHA 2009). These phthalates, together with recently added diisobutyl phthalate (DIBP), have been included into the list of candidates as “substances of very high concern” under the REACH Regulation (ECHA 2010).

Within the EU, DEHP is listed as a priority substance under the Water Framework directive, a regulation designed to improve the quality of water within the EU (EU 2008). DEHP and DnBP have also been identified as substances for priority action under the OSPAR convention, under which signatory countries have agreed a target of cessation of discharges, emissions and losses of all hazardous substances to the marine environment of the North-East Atlantic by 2020, the “one generation” cessation target (OSPAR 1998). In August 2012, despite a European Commission ruling from June 2012 (ENDS 2012), the Danish Ministry of Environment announced plan to introduce a wider ban on marketing and use for four hormone-disrupting phthalates; DEHP, DBP, BBP and DIBP (DMOE 2012).

As far as we are aware, legislation does not currently exist in any of the countries where the 31 tested articles were sold that prohibits the sale of textiles products containing phthalates, although this situation may soon change in Denmark (in which six of the products were purchased for this study), as national prohibitions have recently been announced for four phthalates (DMOE 2012). It is already the case, however, that the use of di-2-ethylhexyl phthalate (DEHP), di-n-butyl phthalate (DnBP) and butyl benzyl phthalate (BBP) are prohibited in all toys or childcare articles put on the market within the EU (with a limit of 0.1% by weight, equivalent to 1000 mg/kg), and the use of diisononyl phthalate (DINP), diisodecyl phthalate (DIDP) and di-n-octyl phthalate (DnOP) is prohibited in such articles if they can be placed in the mouth by children (EU 2005).

For all plastisol print-bearing fabrics that were analysed for phthalate concentrations, a portion of printed fabric was also analysed for the concentration of NPEs, and therefore the concentration of NPEs in these articles are included in Table 6 for comparison. For these four samples, the concentrations of NPEs in three samples were in the lower end of the range of concentrations found across all samples, with NPEs not detected for one sample (TX12008). However, given the relatively small number of samples, it is not possible to determine any clear trends between the presence of phthalates at higher concentrations, and the concentrations of NPEs in plastisol printed fabric.

3.4 Chemical screening

As well as the substances identified in the various items (as outlined in Sections 3.1 – 3.3), one or more additional chemicals were identified in almost all of the items that were investigated using the qualitative chemical screening approach (in 61 of 63 items). A summary of the additional chemicals identified using qualitative screening of extractable organic chemicals in 63 items is given in Table 7, below, followed by some background information on the known uses and any intrinsic hazardous properties associated with these substances. Although it has not been possible to determine the concentrations at which these compounds were present in the samples (as the screening method is necessarily qualitative, not quantitative), the results nevertheless provide an indication of the complexity of chemical residues which can remain in finished textile products, perhaps as a consequence of their use in manufacture, and therefore of the potential for discharge of complex chemical mixtures in effluents generated by textile factories and in laundry wash-waters.

A full listing of all substances identified in individual samples is provided in Appendix 3.

Substance	Number of items (of 63 in total)
Linear alkanes	59
Benzenemethanamine, N-(phenylmethyl)-	1
Benzene, 1,1'-(3-methyl-1-propene-1,3-diyl)bis-	1
Benzophenone	2
Benzyl benzoate	12
1,1'-Biphenyl	2
Benzyl naphthyl ether	1
Bis(2-ethylhexyl) maleate	1
2,6-Di-tert-butyl-4-methyl-phenol (butylated hydroxytoluene, BHT)	2
Ethanol, 2-(2-butoxyethoxy)- / acetate derivative	2
Ethanol, 2-(tetradecyloxy)-	1
Nonylphenol	1
Hexadecanoic acid and esters	8
Octadecanoic acid and esters	5
Octadecenoic acid and esters	2
Octadecadienoic acid and esters	1
Octyl-diphenylamine	2
<i>Others:</i>	
Amyrin	2
α-Amyrenone	20
Sitosterol	21
Cholesterol	3
Squalene	5

Table 7. Substances and substance groups identified by chemical screening, with the number of items in which the substance/group was identified.

Alkanes. Medium and long-chained linear alkanes (from C16 to C36) were commonly identified in the samples investigated, with one or more example being present in 59 of the 63 items tested. These compounds, which are readily biodegradable, are components of petroleum mixtures such as crude oil and refined petroleum products, and some can also originate from plant sources (Ahad et al. 2011, Overton 1994). Longer-chained alkanes are used within textile manufacturing, including as part of finishing processes (Lacasse & Baumann 2004). Some medium and long-chained linear alkanes can have toxicological effects, though mixtures of longer-chain alkanes are used as food additives and these compounds are generally not considered to be of particular toxicological concern in mammals except in high doses (Griffis et al. 2010).

Benzyl benzoate. Other than linear alkanes, benzyl benzoate was the most commonly identified compound, being found in 12 items (see Appendix 3 for details). Benzyl benzoate is a readily biodegradable substance that is used in some dye formulation (IPPC 2003). However, this chemical has been classified under the Globally Harmonised System of Classification and Labelling of Chemicals (GHS)² as toxic to aquatic life with long lasting effects (H411) and harmful if swallowed (H302).

A number of other compounds, or groups of compounds, were identified in two or more of the items tested, including:

Benzophenone (in TX12007 & TX12087), widely used as a fragrance chemical and, occasionally, as a flavouring ingredient. It is also used in the manufacture of insecticides and other agricultural chemicals, and is used as an additive for certain plastics and adhesives (US DHHS 2000). Benzophenone has been classified under GHS as very toxic to aquatic life with long-lasting effects (H410). The liver has been shown to be the primary target organ of benzophenone toxicity in rats and mice, with effects also seen in the kidney for rats (US DHHS 2000).

1,1'-Biphenyl (in TX12015 & TX12121) occurs naturally in petroleum mixtures such as crude oil (Overton 1994), and can also be released through incomplete combustion of petroleum products and coal, including within emissions from vehicles (IPCS 1999). It has been used as a dyestuff carrier for textiles, among other uses (IPCS 1999). This chemical has been classified under the GHS as very toxic to aquatic life with long lasting effects (H410) as well as irritating to eyes and skin (H315, H319).

2,6-Di-tert-butyl-4-methyl-phenol, also known as **butylated hydroxytoluene (BHT)**, (in TX12122 & TX12126), has been widely used as an antioxidant in a wide range of products including plastics and other petrochemical products, cosmetics and food products (Jobling et al. 1995). BHT has been classified under the GHS as toxic to aquatic life with long lasting effects (H411), and irritating to the skin and eyes (H315, H319).

2-(2-Butoxyethoxy)-ethanol, also known as **di-ethylene glycol monobutyl ether (DEGBE)**, was identified in TX12088, and the derivative **2-(2-butoxyethoxy)-ethanol acetate** was identified in TX12127. DEGBE is widely used as a solvent, especially in the printing ink and paint industries, and di-ethylene glycol derivatives in general, including acetates, are used in a variety of textile manufacturing processes and formulations (IPPC 2003). The two compounds that were identified are both are classified under the GHS as irritating to the eyes (H319).

Fatty acids. A number of fatty acids and their ester derivatives were identified relatively frequently, with 12 items containing one or more of the chemicals from this group (see Appendix 3 for details). Hexadecanoic acid (also known as palmitic acid) was identified in two items, and various ester derivatives (methyl, butyl, octadecyl or isopropyl esters) were identified in six other items.

A closely related fatty acid, octadecanoic acid (also known as stearic acid), was identified in one item, with either the methyl or isopropyl ester being identified in four others. The methyl esters of octadecanoic acid (or oleic acid) or octadecadienoic acid (or linoleic acid) were each identified in one other item, and octadecanoic acid itself was identified in another item.

Fatty acid esters, which are present in animal or vegetable fats, oils, or waxes, are used in a variety of textile manufacturing processes, including as defoaming and dispersing agents in dye formulations, and as sizing agents. These, and other fatty acid derivatives such as fatty acid ethoxylates which are used as surfactants, can undergo partial degradation to give rise to the fatty acids themselves (IPPC 2003).

² The Globally Harmonised System of Classification and Labelling of Chemicals (GHS), a system set up under the UN, provides a way of assessing the hazardous properties of chemicals through the use of hazard statements (UN 2011). The hazard statements used for individual substances within Section 3.4 are a composite of information drawn from a number of sources including material safety data sheets (MSDSs) supplied by Sigma-Aldrich (<http://www.sigmaaldrich.com>); Landolt-Börnstein (<http://lb.chemie.uni-hamburg.de>); Merck Millipore (<http://www.millipore.com>); Alfa (<http://www.alfa.com>); ACROS (<http://www.acros.be>)

Fatty acids and their derivatives are not of particular environmental concern due to their low toxicity and being readily biodegradable, and the compounds identified in this study are generally not categorised as hazardous according to the GHS, with the exception of some being able to cause eye irritation (H319), and for some of these also skin irritation (H315).

Other substances were identified in single items amongst the 63 products tested, these included;

Nonylphenol (in TX12127), a persistent, bioaccumulative and toxic substance used to manufacture NPEs, which can degrade in the environment back to nonylphenol. However, NPEs were not detected in this item above the method limit of 1 mg/kg (see Section 3.1). Due to persistence and widespread release, nonylphenol is widely distributed in the environment, having been detected in various media including within surface water environments and in indoor dusts (OSPAR 2004, Butte & Heinzow 2002). See Box A for more information on nonylphenol and NPEs.

Benzenemethanamine, N-(phenylmethyl)-, also known as **N,N-dibenzylamine**, (in TX12063) is a decomposition product of a compound used in rubber vulcanisation processes and has been found in rubber products (Niessner & Klampfl 2000), though it is not clear if the item in which this was identified contained any rubber-based materials. N,N-dibenzylamine has been classified under the GHS as harmful if swallowed (H302) and irritating to eyes and skin (H315, H319).

Benzene, 1,1'-(3-methyl-1-propene-1,3-diyl)bis-, also known as **distyrene**, (in TX12121). It was not possible to identify possible origins of this compound, nor any categorisation according to the GHS

Bis(2-ethylhexyl) maleate, also known as **dioctyl maleate**, (in TX12085) is used as a plasticiser in emulsion-type paints, papers, textile coating, adhesives and oil additives, and is also used in cosmetic skin and hair products, which has resulted in cases of allergic contact dermatitis (Chan & Wakelin 2006, Laguna et al. 2009). Bis(2-ethylhexyl) maleate has been classified under the GHS as irritating to eyes and skin (H315, H319).

Ethanol, 2-(tetradecyloxy)-, also known as **ethylene glycol monotetradecyl ether** or **tetradecylglycol**, (in TX12083), which can be used as a surfactant (Shinoda et al. 1993). As noted above, many glycol derivatives are used in a variety of textile manufacturing processes and formulations (IPPC 2003). It was not possible to identify any categorisation of this glycol derivative according to the GHS, nor other information on its hazardous properties.

Benzyl naphthyl ether, also known as **2-(phenylmethoxy) naphthalene**, (in TX12122), the main use of which appears to be in thermal paper and other heat-sensitive recording materials (Terasaki et al. 2008). Different assessments have classified this ether under the GHS as irritating to the skin and eyes (H315, H319), and as either “very toxic to aquatic life with long-lasting effects” (H410) or “may cause long-lasting harmful effects to aquatic life” (H413).

Octyl-diphenylamine (in TX12056), which can be used as a lubricant (Thompson et al. 2007). It was not possible to identify any categorisation of this compound according to the GHS, nor information on its hazardous properties.

In addition to the industrial chemicals described above, three compounds that belong to a class of naturally occurring terpenoids compounds that can be isolated from plants were commonly identified within the 63 items investigated (Hernández-Vázquez et al. 2010, Mallavadhani et al. 1998):

Sitosterol (in 21 items),

α-Amyrenone (in 20 items),

Amyrin (in 2 items)

Furthermore, two additional naturally occurring compounds were identified in a number of samples; **squalene** (in 5 items) and **cholesterol** (in 3 items). These two compounds have many possible sources, including being part of natural oils that are present on human skin (Lu et al. 2009), which could account for their presence in these cases.

4. Conclusions

This study has demonstrated the presence of a number of different hazardous chemicals within a broad range of textile products of the type included in this study, as either components of materials incorporated within the product or residues remaining from use within manufacturing processes.

Among these, the most commonly detected substances were NPEs, with residues identified in products across all brands and almost all countries of manufacture and countries of purchase included in the study, indicating that the use of NPEs is widespread within the international textile industry, including within supply chains used by several major international clothing brands.

The results for the quantification of NPEs are consistent with those from a previous study carried out by Greenpeace International for 78 textile articles that consisted primarily of sports and recreational clothing, and shoes (Brigden et al. 2012, Greenpeace 2011b). In this earlier study, 67% of the articles tested positive for the presence of NPEs (above 1 mg/kg), with levels ranging from just above 1 mg/kg to 27 000 mg/kg, compared to 63% of articles in the current study, for which levels ranged from just above 1 mg/kg to 45 000 mg/kg (see Table 8).

NPE concentration range (mg/kg)	Current study: No. samples (of 141 analysed)	Current study: % samples (of 141 analysed)	Previous study*: %samples (of 78 analysed)
<1	52	37%	33%
1 – 10	27	19%	22%
>10 – 100	34	24%	27%
>100 – 1 000	16	11%	15%
>1 000	12	9%	3%

Table 8. The number and percentage of samples (of the 141 articles tested) within various NPE concentration ranges, compared to a previous study of 78 textile articles.
* Brigden et al. 2012, Greenpeace 2011b.

In the current study, NPEs were detected in one or more product from 13 of the 18 countries of manufacture. For the five countries of manufacture for which NPEs were not detected, only a small number of articles were tested: Cambodia (2 articles); Jordan (1 article); Romania (1 article); Spain (1 article); and Tunisia (3 articles). Given the low number of articles for these five countries, and the results of the study as a whole, the fact that NPEs were not identified in products manufactured in these countries cannot be taken to indicate that textile products manufactured in these countries in general do not contain detectable residues of NPEs.

The presence of NPEs in finished products indicates their use during manufacture, which can result in releases of NPEs and nonylphenol from manufacturing facilities within the countries of production as well as residues in the products. In addition, NPE residues within textile products are readily released when the items are washed as part of their normal use (Brigden et al. 2012, Greenpeace 2012), resulting in inputs of NPEs and nonylphenol to the environment of the countries in which these products are sold.

The use of NPEs and nonylphenol has effectively been banned within the EU (EU 2003), with similar restrictions in place in the US and Canada (CEPA 2004, USEPA 2010). The manufacture, use and release of NP and NPEs are not currently regulated in China, though NPEs and nonylphenol have recently been included on the “List of toxic chemicals severely restricted for import and export in China”, and their import or export now requires prior permission (MEP 2011). Regulations do not currently exist in EU, or elsewhere, that restrict the sale of textile products containing NPE residues, though such a regulation is currently under development within the EU (KEMI 2012). In order to offer adequate protection, such regulations will need to set any limit for NPEs in products as low as possible and cover as wide a range of NPEs as possible. At the same time, parallel measures to restrict the use of NPEs in manufacture are also needed in countries in which textile products are manufactured.

Some major clothing brands set limits on the presence of certain hazardous substances in their products, as part of their programmes to ensure product safety. However, limits typically set for the presence of NPEs and other alkylphenol ethoxylates (APEs) are currently too high to prevent the continued use of these chemicals during manufacture, and therefore their discharge in the country of manufacture and release through laundering in the country of sale. Details of relevant regulatory measures and the policies of textile brands are discussed in detail in previous reports (Brigden et al. 2012, Greenpeace 2012).

In addition to the presence of NPEs, the identification of phthalates at high concentrations in some articles indicates the ongoing use of phthalates as plasticisers in formulations used to produce plastisol prints on textile products manufactured in a number of countries around the world. The levels and types of individual phthalates found in the four articles with high concentrations (see Table 6) are consistent with previous investigations of related textile products (RAPEX 2012, Greenpeace 2004). The presence of phthalates in textiles, such as through the use as plasticisers in plastisol prints, is not currently subject to any regulations in the countries where the items were sold, though further national controls are pending in Denmark. However, regulations do exist within the EU that prohibit phthalates (including those identified in this study) in toys and childcare articles (EU 2005), thereby recognising the hazard associated with products containing phthalates, including plastic parts containing phthalates as plasticisers. Regulations and brand policies are required to address the hazard that the presence of phthalates poses in a broad range of products, including textiles items.

Carcinogenic amines released under reducing conditions were detected in only a small fraction of items tested (2 of 134 items). Previous studies have similarly identified carcinogenic amines for a relatively low fraction of items tested (JRC 2008, Laursen et al. 2003), including items for which far higher concentrations of amines were detected compared to levels found in this study. For each of the two items in this study in which amines were detected, the concentration of the amine (*o*-dianisidine) was below limits set by regulations that address the release of carcinogenic amines from textiles treated with azo dyes. For these items, the source of *o*-dianisidine at the levels found may have been the presence of an azo dye manufactured from *o*-dianisidine. However, residues of this amine within other types of azo dye formulations used during manufacture may have contributed to the *o*-dianisidine identified. Whatever the source or level identified, the presence of any substances in textiles capable of releasing a carcinogenic amine following reduction is clearly cause for concern.

In addition to the chemicals that were quantified in the textile products, a wide range of other chemicals was identified in various items through qualitative chemical screening. While not all the chemicals identified are known to have hazardous properties, many have been classified under the GHS as having hazardous properties, including some being classified as being very toxic to aquatic life with long-lasting effects (H410). Many of the chemicals that were identified have known uses within textile manufacturing, though some do also have other potential sources unrelated to use during textile manufacture. In addition, three compounds of plant origin and two compounds that can originate from natural oils present on human skin were identified in many of the items investigated. In most cases, the industrial chemicals identified through screening were found in only one or two of the 63 items investigated in this way, though certain chemicals or chemicals groups were more commonly found, particularly linear alkanes and benzyl benzoate, as well as fatty acids and their derivatives. For the items investigated, there were no clear associations within individual samples in terms of chemicals identified, other than for fatty acids and their esters to some extent. That is, for each sample, there were no clear patterns in which the presence of one particular chemical was commonly accompanied by the presence of another, with certain exceptions for fatty acids and their esters. For two samples (TX12029 & TX12123), the methyl esters of both hexadecanoic acid and octadecanoic acid were identified in each sample, and similarly both hexadecanoic acid and octadecanoic acid were identified in one other sample (TX12127). These results indicate there may be a common origin for these fatty acids and their methyl esters in some cases.

A number of the compounds identified in this study using a qualitative chemical screening approach have previously been identified in textile items sold in Denmark that were investigated using a similar screening approach, including alkanes, benzyl benzoate, fatty acids, bis(2-ethylhexyl) maleate and squalene (Laursen et al. 2003). Also, glycols related to those identified in the current study were identified in this previous Danish study. The screening aspect of the study highlights the diverse range of chemical residues that can be present within textile products, though from the results of this study do not reveal any associations between the presence of individual chemicals identified through the screening approach and factors such as the type of item, where it was manufactured or sold, or the item brand. However, the qualitative screening data do suggest that analyses that focus only on a narrow range of target substances will underestimate the complexity of chemical residues in finished textiles and, therefore, the potential for chemical releases during manufacture and subsequent laundering.

Overall, this study has provided a greater understanding of the presence and, in some cases, concentrations of a broad range of chemicals within high street fashion textile products, across a diverse range of brands and countries of manufacture and sale. Despite the scale of this study, the number of articles investigated is, of course, inevitably small compared to the vast number of products manufactured and sold per country or per brand. It cannot, therefore, be assumed that the results obtained in this study are representative of levels or presence of the chemicals investigated which may be expected for all such products. Rather, this study provides a snapshot of what appears to be a more generic problem that is not restricted to any particular country, product type or brand, and one that deserves further investigation including from a regulatory and brand policy perspective.

5. References

- Abb M, Heinrich T, Sorkau E & Lorenz W (2009).** Phthalates in house dust. *Environment International* 35(6): 965-970
- Abdul-Ghani S, Yanai J, Abdul-Ghani R, Pinkas A & Abdeen Z (2012).** The teratogenicity and behavioral teratogenicity of di(2-ethylhexyl) phthalate (DEHP) and di-butyl phthalate (DBP) in a chick model. *Neurotoxicology and Teratology* 34(1): 56-62
- Ahad JME, Ganeshram RS, Bryant CL, Cisneros-Dozal LM, Ascough PL, Fallick AE, Slater GF (2011).** Sources of n-alkanes in an urbanized estuary: Insights from molecular distributions and compound-specific stable and radiocarbon isotopes *Marine Chemistry* 126(1–4): 239-249
- Aso S, Ehara H, Miyata K, Hosyuyama S, Shiraishi K, Umano T & Minobe Y (2005).** A two-generation reproductive toxicity study of butyl benzyl phthalate in rats. *Journal of Toxicological Sciences* 30(SI): 39-58
- Blount BC, Silva MJ, Caudill SP, Needham LL, Pirkle JL, Sampson EJ, Lucier GW, Jackson RJ & Brock JW (2000).** Levels of seven urinary phthalate metabolites in a human reference population. *Environmental Health Perspectives* 108(10): 979-982
- Boberg J, Christiansen S, Axelstad M, Kledal TS, Vinggaard AM, Dalgaard M, Nellemann C & Hass U (2011).** Reproductive and behavioral effects of di-isononyl phthalate (DiNP) in perinatally exposed rats. *Reproductive Toxicology* 31(2): 200-209
- Brennan RJ & Schiestl RH (1999).** The aromatic amine carcinogens *o*-toluidine and *o*-anisidine induce free radicals and intrachromosomal recombination in *Saccharomyces cerevisiae*. *Mutation Research* 430: 37-45
- Brigden K, Santillo D & Johnston P (2012).** Nonylphenol ethoxylates (NPEs) in textile products, and their release through laundering. Greenpeace Research Laboratories Technical Report 01/2012, 14pp.
http://www.greenpeace.to/greenpeace/wp-content/uploads/2012/03/Dirty_Laundry_Product_Testing_Technical_Report_01-2012.pdf
- Butte W & Heinzow B (2002).** Pollutants in house dust as indicators of indoor contamination. *Reviews in Environmental Contamination and Toxicology* 175: 1-46
- Cadogan DF, Papez M, Poppe AC, Pugh DM & Scheubel J (1993).** In: *PVC 93: The Future*. Publ: The Institute of Materials; pp260-274
- Caldwell JC (2012).** DEHP: Genotoxicity and potential carcinogenic mechanisms—A review. *Mutation Research/Reviews in Mutation Research*, In Press, Corrected Proof, Available online 3 April 2012
- CEPA (2004).** Notice requiring the preparation and implementation of pollution prevention plans in respect of effluents from textile mills that use wet processing (TMEs) and nonylphenol (NP) and its ethoxylates (NPEs), under the Canadian Environmental Protection Act (CEPA), 1999. *Canada Gazette Part I, Vol. 138, No. 49, 4th December 2004*. <http://www.ec.gc.ca/planp2-p2plan/B2D19B6D-325F-458A-88E1-F69291E58DE3/g1-13849.pdf>
- Colon I, Caro D, Bourdony CJ & Rosario O (2000).** Identification of phthalate esters in the serum of young Puerto Rican girls with premature breast development. *Environmental Health Perspectives* 108(9): 895-900
- Dalgaard M, Nellemann C, Lam HR, Sorensen IK & Ladefoged O (2001).** The acute effects of mono(2-ethylhexyl)phthalate (MEHP) on testes of prepubertal Wistar rats. *Toxicology Letters* 122: 69-79
- DHHS (2011).** 3,3'-Dimethoxybenzidine and dyes metabolized to 3,3'-dimethoxybenzidine. Report on carcinogens, 12th Edition. U.S. Department of Health and Human Service. Public Health Service National Toxicology Program. <http://ntp.niehs.nih.gov/ntp/roc/twelfth/roc12.pdf>
- DMOE (2012).** 'Danish Ministry of the Environment protects consumers from dangerous phthalates' announcement by the Danish Ministry of the Environment (DMOE), 23rd August 2012.
http://www.mim.dk/Nyheder/20120823_ftalater.htm (in Danish)
- DoE (1991).** Environmental hazard assessment: Di(2-ethylhexyl)phthalate. Report TSD/2 Publ: United Kingdom Department of the Environment Toxic Substances Division. 51pp.
- Dunkel VCE, Zeiger D, Brusick E, McCoy D, McGregor K, Mortelmans HS, Rosenkranz & Simmon VF (1985).** Reproducibility of microbial assays, II. Testing of carcinogens and noncarcinogens in *Salmonella typhimurium* and *Escherichia coli*, *Environ. Mutagen.*, 7 (Suppl. 5): 1-248
- EC (2005).** Directive 2005/84/EC of the European Parliament and of the Council of 14 December 2005 amending for the 22nd time Council Directive 76/769/EEC on the approximation of the laws, regulations and administrative

provisions of the Member States relating to restrictions on the marketing and use of certain dangerous substances and preparations (phthalates in toys and childcare articles). Official Journal of the European Communities L344, 27.12.2005: 40-43 <http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2005:344:0040:0043:EN>

ECB (2002). European Union Risk Assessment for o-Anisidine (CAS No 90-04-0), Final Report p.50. European Chemicals Bureau (ECB)

ECHA (2010). Candidate List of Substances of Very High Concern for Authorisation, publ. European Chemicals Agency (ECHA), 13.01.2010
http://www.precidip.com/data/files/pdf/Candidate_List_of_Substances_of_Very_High_Concern_for_authorisation.pdf
(accessed 23.08.2012)

ECHA (2011). Candidate List of Substances of Very High Concern for authorization. European Chemicals Agency.
http://echa.europa.eu/chem_data/authorisation_process/candidate_list_table_en.asp

Ema M & Miyawaki E (2002). Effects on development of the reproductive system in male offspring of rats given butyl benzyl phthalate during late pregnancy. Reproductive Toxicology 16: 71-76

ENDS (2012). Danish Phthalate ban unnecessary – experts. <http://www.endseurope.com/29054/danish-phthalate-ban-unnecessary-experts> accessed 23.08.2012

EU (2002). Directive 2002/61/EC of the European Parliament and of the Council of 19 July 2002 amending for the 19th time Council Directive 76/769/EEC relating to restrictions on the marketing and use of certain dangerous substances and preparations (azocolourants): [<http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2002:243:0015:0018:EN:PDF>]

EU (2003). Directive 2003/53/EC Of The European Parliament And Of The Council Of 18 June 2003 Amending For The 26th Time Council Directive 76/769/EEC Relating To restrictions on the marketing and use of certain dangerous substances and preparations (nonylphenol, nonylphenol ethoxylate and cement)
http://europa.eu.int/eur-lex/pri/en/oj/dat/2003/l_178/l_17820030717en00240027.pdf

EU (2005). Directive 2005/84/EC of the European Parliament and of the Council of 14 December 2005 amending for the 22nd time Council Directive 76/769/EEC on the approximation of the laws, regulations and administrative provisions of the Member States relating to restrictions on the marketing and use of certain dangerous substances and preparations (phthalates in toys and childcare articles): [<http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2005:344:0040:0043:EN:PDF>]

EU (2008). Directive 2008/105/EC of the European Parliament and of the Council of 16 December 2008 on environmental quality standards in the field of water policy, amending and subsequently repealing Council Directives 82/176/EEC, 83/513/EEC, 84/156/EEC, 84/491/EEC, 86/280/EEC and amending Directive 2000/60/EC of the European Parliament and of the Council. Official Journal of the European Union L 348:84-97

Fasano E, Bono-Blay F, Cirillo T, Montuori P & Lacorte S (2012). Migration of phthalates, alkylphenols, bisphenol A and di(2-ethylhexyl)adipate from food packaging. Food Control 27(1): 132-138

Ferri M, Chiellini F, Pili G, Grimaldi L, Florio ET, Pili S, Cucci F & Latini G (2012). Di-(2-ethylhexyl)-phthalate migration from irradiated poly(vinyl chloride) blood bags for graft-vs-host disease prevention. International Journal of Pharmaceutics 430(1–2):Pages 86-88

Fierens T, Servaes K, Van Holderbeke M, Geerts L, De Henauf S, Sioen I & Vanermen G (2012). Analysis of phthalates in food products and packaging materials sold on the Belgian market. Food and Chemical Toxicology 50(7): 2575-2583

Fromme H, Lahrz T, Piloty M, Gebhart H, Oddoy A & Rüdén H (2004). Occurrence of phthalates and musk fragrances in indoor air and dust from apartments and kindergartens in Berlin (Germany). Indoor Air 14 (3): 188-195

Golka K, Kopps S, Myslak ZW (2004). Carcinogenicity of azo colorants: influence of solubility and bioavailability. Toxicology Letters 151(1): 203-210

Grande SW, Andrade AJ, Talsness CE, Grote K & Chahoud I (2006). A dose–response study following in utero and lactational exposure to di(2-ethylhexyl)phthalate: effects on female rat reproductive development. Toxicol. Sci. 91: 247–254

Gray LE (Jnr), Laskey J & Ostby J (2006). Chronic di-n-butyl phthalate exposure in rats reduces fertility and alters ovarian function during pregnancy in female Long Evans hooded rats. Toxicol. Sci. 93: 189–195

Greenpeace (2004). Finding Chemo - Toxic Childrenswear by Disney. Greenpeace International, pp18.
<http://www.greenpeace.org/international/en/publications/reports/finding-chemo-toxic-children/>

Greenpeace (2011a). Dirty Laundry. Unravelling the corporate connections to toxic water pollution in China, pp 116; including the accompanying Technical Note, pp55.
<http://www.greenpeace.org/dirtylaundryreport>

Greenpeace (2011b). Dirty Laundry 2: Hung Out to Dry - Unravelling the toxic trail from pipes to products, pp32.
<http://www.greenpeace.org/international/en/publications/reports/Dirty-Laundry-2/>

Greenpeace (2012). Dirty Laundry: Reloaded - How big brands are making consumers unwitting accomplices in the toxic water cycle, pp48. <http://www.greenpeace.org/international/en/publications/Campaign-reports/Toxics-reports/Dirty-Laundry-Reloaded/>

Griffis LC, Twerdok LE, Francke-Carroll S, Biles RW, Schroeder RE, Bolte H, Faust H, Hall WC & Rojko J (2010). Comparative 90-day dietary study of paraffin wax in Fischer-344 and Sprague–Dawley rats. Food and Chemical Toxicology 48(1): 363-372

Guerranti C, Sbordoni I, Fanello EL, Borghini F, Corsi I & Focardi SI (2012). Levels of phthalates in human milk samples from central Italy. Microchemical Journal, In Press, Corrected Proof, Available online 8 July 2012

Haley TJ (1975). Benzidine revisited: A review of the literature and problems associated with the use of benzidine and its congeners. Clinical Toxicology 8(1): 13-42

Hernández-Vázquez L, Mangas S, Palazón J & Navarro-Ocaña A (2010). Valuable medicinal plants and resins: Commercial phytochemicals with bioactive properties. Industrial Crops and Products 31(3): 476-480

Howdeshell KL, Wilson VS, Furr J, Lambright CR, Rider CV, Blystone CR, Hotchkiss AK & Gray LE (Jnr) (2008). A mixture of five phthalate esters inhibits fetal testicular testosterone production in the Sprague Dawley rat in a cumulative dose additive manner. Toxicol. Sci. 105: 153–165

IARC (1987). Overall Evaluations of Carcinogenicity: An Updating of IARC Monographs Volumes 1 to 42, supplement 7. International Agency for Research on Cancer (IARC).
<http://monographs.iarc.fr/ENG/Monographs/suppl7/index.php>

IARC (1998). Aromatic amines. In: International Agency for Research on Cancer (IARC) monographs on the evaluation of the carcinogenic risk of chemicals to humans. Volume 4; Some aromatic amines, hydrazine and related substances, N-nitroso compounds and miscellaneous alkylating agents, updated 1998.
<http://monographs.iarc.fr/ENG/Monographs/vol4/volume4.pdf>

IARC (2008). International Agency for Research on Cancer (IARC) monographs on the evaluation of the carcinogenic risk of chemicals to humans. Volume 99; Some Aromatic Amines, Organic Dyes, and Related Exposures. <http://monographs.iarc.fr/ENG/Monographs/vol99/mono99.pdf>

IPCS (1999). Biphenyl. Concise international chemical assessment document: 6. International Programme on Chemical Safety (IPCS). ISBN 92-4-153006-5, Geneva 1999

IPPC (2003). Reference document on best available techniques for the textiles industry, Integrated Pollution Prevention and Control (IPPC), European Commission

Jenke DR, Story J & Lalani R (2006). Extractables/leachables from plastic tubing used in product manufacturing. International Journal of Pharmaceutics 315(1–2): 75-92

Jobling S, Sheahan D, Osborne JA, Matthiessen P & Sumpter JP (1996). Inhibition of testicular growth in rainbow trout (*Oncorhynchus mykiss*) exposed to estrogenic alkylphenolic chemicals. Environmental Toxicology and Chemistry 15(2): 194-202

Jobling S, Reynolds T, White R, Parker MG & Sumpter JP (1995). A variety of environmentally persistent chemicals, including some phthalate plasticizers, are weakly estrogenic. Environmental Health Perspectives 103(6): 582-587

JRC (2008). European survey on the presence of banned azodyes in textiles, EUR 23447 EN – 2008. Joint Research Commission, Institute for Health and Consumer Protection, European Commission. ISBN 978-92-79-09118-6. http://publications.jrc.ec.europa.eu/repository/bitstream/111111111/1321/1/eur_23447_en_fr_aa.pdf

KEMI (2012). Proposals for new restrictions under REACH. Swedish Chemicals Agency (KEMI).
<http://www.kemi.se/en/Content/Rules-and-regulations/Reach/Begransningsregler-bilaga-XVII/Proposals-for-new-restrictions/>

- Kirk-Othmer (1994).** Encyclopedia of Chemical Technology. 4th ed. Volume 1: New York, NY. John Wiley and Sons, 1991-Present, V9, p. 1062
- Koch HM, Preuss R & Angerer J (2006).** Di(2-ethylhexyl)phthalate (DEHP): human metabolism and internal exposure—an update and latest results. *Int. J. Androl.* 29: 155–165
- Lacasse K & Baumann W (2004).** Textile Chemicals: Environmental data and facts. Springer- Verlag ISBN 3-540-40815-0
- Laguna C, de la Cuadra J, Martín-González B, Zaragoza V, Martínez-Casimiro L & Alegre V (2009).** Allergic Contact Dermatitis to Cosmetics. *Actas Dermo-Sifiliográficas (English Edition)*, 100(1): 53-60
- Langer S, Weschler CJ, Fischer A, Bekö G, Toftum L & Clausen G (2010).** Phthalate and PAH concentrations in dust collected from Danish homes and daycare centers. *Atmospheric Environment* 44(19):2294-2301
- Latorre I, Hwang S, Sevillano M & Montalvo-Rodríguez R (2012).** PVC biodeterioration and DEHP leaching by DEHP-degrading bacteria. *International Biodeterioration & Biodegradation* 69:73-81
- Laursen SE, Hansen J, Drøjdahl A, Hansen OC, Pommer K, Pedersen E & Bernth N (2003).** Survey of chemical compounds in textile fabrics. Survey no. 23, on behalf of the Danish Environmental Protection Agency. <http://www.mst.dk/NR/rdonlyres/B9CDE217-9E41-4F27-A8A3-921D5B50A737/0/23.pdf>
- Liao Ch-S, Yen J-H & Wang Y-Sh (2006).** Effects of endocrine disruptor di-n-butyl phthalate on the growth of Bok choy (*Brassica rapa* subsp. *chinensis*). *Chemosphere* 65: 1715–172
- Lin H, Ge R-S, Chen G-R, Hu G-X, Dong L, Lian QQ, Hardy DO, Sottas CM, Li X-K & Hardy MP (2008).** Involvement of testicular growth factors in fetal Leydig cell aggregation after exposure to phthalate in utero. *Proc. Natl Acad. Sci. USA* 105(20): 7218–7222
- Liu H, Liang Y, Zhang D, Wang C, Liang H & Cai H (2010).** Impact of MSW landfill on the environmental contamination of phthalate esters. *Waste Management* 30(8–9):1569-1576
- Lovekamp-Swan T & Davis BJ (2003).** Mechanisms of phthalate ester toxicity in the female reproductive system. *Environmental Health Perspectives* 111(2): 139-145
- Lu GW, Valiveti S, Spence J, Zhuang C, Robosky L, Wade K, Love A, Hu L-Y, Pole D, Mollan M (2009).** Comparison of artificial sebum with human and hamster sebum samples. *International Journal of Pharmaceutics*, Vol 367(1–2): 37-43
- Mallavadhani UV, Panda AK & Rao YR (1998).** Pharmacology and chemotaxonomy of diospyros, review article number 134. *Phytochemistry* 49(4): 901-951
- Martino-Andrade AJ, Morais RN, Botelho GG, Muller G, Grande SW, Carpentieri GB, Leão GM & Dalsenter PR (2008).** Coadministration of active phthalates results in disruption of foetal testicular function in rats. *Int. J. Androl.*, Dec. 2008: 9pp. (DOI 10.1111/j.1365-2605.2008.00939)
- MEP (2011).** List of Toxic Chemicals Severely Restricted on the Import and Export in China. Ministry of Environmental Protection (MEP). http://www.crc-mep.org.cn/news/NEWS_DP.aspx?TitID=267&T0=10000&LanguageType=CH&Sub=125 (in Chinese)
- Morgan DL, Dunnick JK, Goehl T, Jokinen MP, Matthews HB, Zeiger E & Mennear JH (1994).** Summary of the National Toxicology Program Benzidine Dye Initiative. *Environmental Health Perspectives* 102(suppl 12): 63-78
- Mylchreest E, Sar M, Wallace DG & Foster PMD (2002).** Fetal testosterone insufficiency and abnormal proliferation of Leydig cells and gonocytes in rats exposed to di(n-butyl) phthalate. *Reproductive Toxicology* 16: 19-28
- Niessner G & Klampfl CW (2000).** Direct comparison of solid-phase extraction and solid-phase microextraction for the gas chromatographic determination of dibenzylamine in artificial saliva leachates from baby bottle teats. *Analytica Chimica Acta* 414(1–2): 133-140
- OSPAR (1998).** OSPAR Strategy with Regard to Hazardous Substances, OSPAR Convention for the Protection of the Marine Environment of the North-East Atlantic, OSPAR 98/14/1 Annex 34
- OSPAR (2004).** Nonylphenol/nonylphenoethoxylates, OSPAR Priority Substances Series 2001, updated 2004, OSPAR Convention for the Protection of the Marine Environment of the North-East Atlantic, OSPAR Commission, London, ISBN 0-946956-79-0: 20 pp.

- Otake T, Yoshinaga J & Yanagisawa Y (2001).** Analysis of organic esters of plasticizer in indoor air by GC-MS and GC-FPD. *Environmental Science and Technology* 35(15): 3099-3102
- Overton EB (1994).** Toxicity of petroleum. In: *Basic Environmental Toxicology*. Cockerham & Shane [Eds], Chapter 5: 133-156
- Rafi F, Hall JD & Cerniglia CE (1997).** Mutagenicity of azo dyes used in foods, drugs and cosmetics before and after reduction by *Clostridium* species from the human intestinal tract. *Food and Chemical Toxicology* 35(9): 897-901
- RAPEX (2012).** RAPEX notifications, European Commission.
http://ec.europa.eu/consumers/dyna/rapex/rapex_archives_en.cfm
- SAPRC (2012).** GB18401-2010, National general safety technical code for textile products. General Administration of Quality Supervision, Inspection and Quarantine of the People's Republic of China, Standardization Administration of the People's Republic of China (SAPRC)
- Sasaki YF, Nishidate E, Su YQ, Matsusaka N, Tsuda S, Susa N, Furukawa Y & Ueno S (1998).** Organ-specific genotoxicity of the potent rodent bladder carcinogens *o*-anisidine and *p*-cresidine. *Mutation Research* 412(2): 155-60
- Shinoda K, Yamaguchi N & Usui M (1993).** Critical composition in organized surfactant solutions. *Colloids and Surfaces A: Physicochemical and Engineering Aspects* 79(2-3): 151-155
- Silva MJ, Barr DB, Reidy JA, Malek NA, Hodge CC, Caudill SP, Brock JW, Needham LL & Calafat AM (2004).** Urinary levels of seven phthalate metabolites in the U.S. population from the National Health and Nutrition Examination Survey (NHANES) 1999-2000. *Environmental Health Perspectives* 112(3): 331-338
- SSNC (2006).** Fluorinated pollutants in all-weather clothing. Swedish Society for Nature Conservation (SSNC).
http://naturvern.imaker.no/data/f/0/75/41/1_2401_0/2_Fluorinated_pollutants_in_all-weather_clothing.pdf
- Swan SH, Main KM, Liu F, Stewart SL, Kruse RL, Calafat AM, Mao CS, Redmon JB, Ternand CL, Sullivan S & Teague JL (2005).** Decrease in anogenital distance among male infants with prenatal phthalate exposure. *Environmental Health Perspectives* 113(8): 1056-1061
- Terasaki M, Fukazawa H, Tani Y & Makino M (2008).** Organic pollutants in paper-recycling process water discharge areas: First detection and emission in aquatic environment. *Environmental Pollution*, 151(1): 53-59
- Thompson BAW, Goldsworthy PM, Riddle MJ, Snape I, Stark JS (2007).** Contamination effects by a 'conventional' and a 'biodegradable' lubricant oil on infaunal recruitment to Antarctic sediments: A field experiment *Journal of Experimental Marine Biology and Ecology* 340(2): 213-226
- Thompson DC, Josephy PD, Chu JWK & Eling TE (1992).** Enhanced mutagenicity of anisidine isomers in bacterial strains containing elevated N-acetyltransferase activity. *Mutation Research* 279: 83-89
- UN (2011).** Globally Harmonised System of Classification and Labelling of Chemicals (GHS), 4th revised edition.
http://www.unece.org/trans/danger/publi/ghs/ghs_rev04/04files_e.html
- US DHHS (2000).** NTP Technical Report on the Toxicity Studies of Benzophenone Administered in Feed to F344/N Rats and B6C3F1 Mice. Toxicity Report Series Number 61, NIH Publication No. 00-3943, pp.8
- USPEA (2010).** Nonylphenol (NP) and Nonylphenol Ethoxylates (NPEs) Action Plan. United States Environmental Protection Agency (USEPA), August 18, 2010. <http://www.epa.gov/oppt/existingchemicals/pubs/actionplans/np-npe.html>
- Verschuieren K (2001).** Handbook of Environmental Data on Organic Chemicals. Volumes 1-2. 4th ed. John Wiley & Sons. New York, NY. p. V1 210

Table A1: Concentrations of NPEs, carcinogenic amines and phthalates in all articles tested.

Details of all articles, including the concentrations of NPEs, carcinogenic amines and phthalates. For NPEs, * indicates the analysis of a section of fabric bearing a plastisol print; for carcinogenic amines “<5 mg/kg” indicates that all quantified amines were below the detection limit (<5 mg/kg) and where a specific amine is listed, all other quantified amines in that sample were below the detection limit (<5 mg/kg); For phthalates, the total concentration of the 9 quantified phthalates is given (mg/kg), with the individual phthalate concentrations provided in Appendix 2; “—” indicates not tested, either due to being undyed fabric (for carcinogenic amines) or article without a medium/large sized plastisol print (for phthalates). TX12066 was not tested as the item was identical to TX12068.

Sample code	Brand	Place of sale	Place of manufacture	Kind of product	Man Woman Child	Fabric	NPEs (mg/kg)	Amines (mg/kg)	Phthalates, total (mg/kg)
TX12001	Armani	Thailand	Mainland China	underwear	woman	96% polyamide, 4% elastane	32	<5	-
TX12002	Armani	Sweden	Mainland China	shirt	---	unknown	43	<5	-
TX12003	Armani	Switzerland	Mainland China	jeans	man	100% cotton excluded decorations	<1	<5	-
TX12004	Armani	UK	Vietnam	polo shirt	man	98% cotton, 2 % elastane	4.8	<5	-
TX12005	Armani	USA	Indonesia	underwear	woman	85% polyamide, 16% elastane	1.2	<5	-
TX12006	Armani	Russia	Thailand	underwear	man	100% cotton	<1	<5	-
TX12007	Armani	Italy	Vietnam	underwear	woman	90% cotton, 10% elastane	<1	<5	-
TX12008	Armani	Italy	Turkey	t-shirt	man	100% cotton	* <1	<5	223 440
TX12009	Armani	France	Indonesia	bra	woman	87% nylon, 13% elastane	8.1	<5	-
TX12010	Benetton	Mexico	Romania	t-shirt	child	100% cotton	* <1	<5	128
TX12011	Benetton	Czech Republic	Tunisia	t-shirt	child	100% organic cotton	* <1	-	33
TX12012	Benetton	Switzerland	Cambodia	hoodie	man	100% cotton	<1	<5	-
TX12013	Benetton	UK	Bangladesh	t-shirt	child	100 % cotton	* <1	-	47
TX12014	Benetton	Russia	Cambodia	t-shirt	child	100% cotton	<1	<5	-
TX12015	Benetton	Italy	Mainland China	jacket	child	outside: 70% cotton, 30% polyamide Inside: 100% polyester	95	<5	-
TX12016	Benetton	Italy	Egypt	sweatshirt	child	100% cotton	11	<5	-
TX12017	Benetton	Belgium	Bangladesh	t-shirt (part of a set)	child	100% cotton	<1	<5	-
TX12018	Benetton	France	India	trousers	child	100% cotton	6.3	<5	-
TX12019	Vero Moda	Mainland China	unknown	top	woman	61% cotton, 37% polyester,2% elastane	31	<5	-
TX12020	Vero Moda	Mainland China	unknown	top	woman	100% polyester	6.3	<5	-
TX12021	Vero Moda	Denmark	India	blouse	woman	unknown	45	<5	-
TX12022	Vero Moda	Denmark	India	top	woman	100% cotton	<1	<5	-
TX12023	Vero Moda	Lebanon	India	dress	woman	100% polyester	130	<5	-
TX12024	Only	Mainland China	unknown	jeans	woman	99% cotton, 1% elastane	5.5	<5	-
TX12025	Only	Mainland China	unknown	t-shirt	woman	Fabric 1- 86% viscose rayon, 5% elastane; Fabric 2-62% viscose rayon, 38% polyester	* 32	-	18
TX12026	Only	Denmark	Mainland China	jeans	woman	unknown	730	<5	-
TX12027	Only	Norway	Turkey	jeans	woman	unknown	38	<5	-
TX12028	Jack & Jones	Mainland China	unknown	t-shirt	man	95% cotton, 5% elastane	* <1	<5	14
TX12029	Jack & Jones	Mainland China	unknown	underwear	man	96% cotton, 4% elastane	2 100	<5	-
TX12030	Jack & Jones	Netherlands	Bangladesh	polo shirt	man	100% cotton	<1	<5	-
TX12031	Jack & Jones	Denmark	Turkey	jeans	woman	unknown	17	<5	-
TX12032	Jack & Jones	Lebanon	Bangladesh	t-shirt	man	85% cotton, 15% viscose	* 4.6	<5	17
TX12033	Calvin Klein	Germany	Egypt	underwear	man	82% polyester, 18% elastane	9.0	<5	-
TX12034	Calvin Klein	Philippines	Mainland China	underwear	man	95% cotton, 5% elastane	20	<5	-
TX12035	Calvin Klein	Mexico	Mexico	jeans	man	100% cotton	56	<5	-
TX12036	Calvin Klein	Netherlands	Jordan	underwear	woman	95% cotton, 5% elastane	<1	<5	-
TX12037	Calvin Klein	USA	Vietnam	jeans	man	100% cotton exclusive of decoration	73	<5	-

Sample code	Brand	Place of sale	Place of manufacture	Kind of product	Man Woman Child	Fabric	NPEs (mg/kg)	Amines (mg/kg)	Phthalates, total (mg/kg)
TX12038	Calvin Klein	South Africa	Thailand	underwear	man	100% cotton	14	<5	-
TX12039	Calvin Klein	Indonesia	Vietnam	underwear	man	92% cotton; 8% Lycra	5.6	<5	-
TX12040	Calvin Klein	Canada	India	t-shirt	man	100% cotton	* 4 000	<5	9
TX12041	C&A	Mexico	Mexico	t-shirt	man	100% cotton	* 45 000	<5	61
TX12042	C&A	Switzerland	unknown	top	child	100% cotton	6.9	-	-
TX12043	C&A	Switzerland	unknown	jacket	child	96% polyester, 4% elastane	64	<5	-
TX12044	C&A	Belgium	unknown	t-shirt	man	100% cotton	* <1	<5	33
TX12045	C&A	Hungary	unknown	t-shirt	child	100% cotton	* 1.7	<5	18
TX12046	C&A	France	unknown	trousers	child	70% cotton, 30% polyester	63	<5	-
TX12047	Diesel	Germany	Morocco	jeans	man	98% cotton, 2% polyurethane	710	<5	-
TX12048	Diesel	Czech Republic	Turkey	vest top	woman	100% cotton	<1	<5	-
TX12049	Diesel	Spain	Mainland China	shorts	man	100% cotton	<1	<5	-
TX12050	Diesel	Russia	Mainland China	t-shirt	man	100% cotton	* 6.6	<5	56
TX12051	Diesel	Italy	Tunisia	jeans	man	100% cotton	<1	<5	-
TX12052	Diesel	Austria	Tunisia	trousers	woman	76% cotton, 22% polyester, 2% elastane	<1	<5	-
TX12053	Diesel	South Africa	India	t-shirt	man	100% cotton	<1	<5	-
TX12054	Diesel	Israel	Mainland China	t-shirt	man	100% cotton	* 16	<5	83
TX12055	Diesel	Hungary	Mainland China	t-shirt	man	100% cotton	* <1	<5	57
TX12056	Esprit	Mainland China	Mainland China	bra	woman	surface 90% cotton, 10% elastane; inside - 100% polyester	<1	<5	-
TX12057	Esprit	Hong Kong	Mainland China	dress	woman	shell: 96% polyester, 4% elastane; lining: 100% polyester	<1	<5	-
TX12058	Esprit	Germany	unknown	t-shirt	youth	100% cotton	* 770	-	14
TX12059	Esprit	Thailand	Mainland China	jacket	woman	100% cotton	460	<5	-
TX12060	Esprit	Finland	Turkey	t-shirt	woman	unknown	<1	<5	-
TX12061	Esprit	Switzerland	unknown	dress	woman	100% cotton	1.1	-	-
TX12062	Esprit	Russia	Mainland China	coat	woman	unknown	17	<5	-
TX12063	Esprit	Belgium	unknown	t-shirt	child	100% cotton	27	<5	-
TX12064	Esprit	Indonesia	Indonesia	dress	woman	outer 100% polyester; inside 100% nylon	66	<5	-
TX12065	Gap	Thailand	Indonesia	jeans	child	99% cotton, 1% elastane	<1	<5	-
TX12066	Gap	Philippines	Indonesia	beach shirt	child	body: 80% polyester, 20% elastane. sleeve: 80% nylon, 20% elastane	-	-	-
TX12067	Gap	Mexico	Mexico	jeans	man	70% cotton, 27% polyester, 3% elastane	920	<5	-
TX12068	Gap	USA	Indonesia	beach shirt	child	body: 80% polyester, 20% elastane. sleeve: 80% nylon, 20% elastane	* <1	<5	14
TX12069	Gap	South Africa	Bangladesh	trousers	man	100% cotton	1.3	<5	-
TX12070	Gap	South Africa	Vietnam	raincoat	child	100% cotton	* 700	<5	14
TX12071	Gap	Israel	Vietnam	dress	woman	100% polyester	43	<5	-
TX12072	Gap	France	Vietnam	t-shirt	child	100% cotton	* 110	<5	25

Sample code	Brand	Place of sale	Place of manufacture	Kind of product	Man Woman Child	Fabric	NPEs (mg/kg)	Amines (mg/kg)	Phthalates, total (mg/kg)
TX12073	Gap	Indonesia	Pakistan	jeans	child	100% cotton	3.8	<5	-
TX12074	Gap	Canada	Indonesia	t-shirt	child	80% nylon 20% elastane	* 8.6	<5	26
TX12075	H&M	Denmark	Turkey	t-shirt	child	unknown	* <1	<5	23
TX12076	H&M	Spain	India	dress	woman	100% polyester	8.7	<5	-
TX12077	H&M	Belgium	Mainland China	sweater	man	100% cotton	<1	<5	-
TX12078	H&M	Lebanon	Mainland China	top	woman	100% polyester	1.6	<5	-
TX12079	H&M	Hungary	Bangladesh	underwear	woman	86% polyamide, 14% elastane	<1	<5	-
TX12080	H&M	France	Mainland China	trousers	child	85% cotton, 14% polyester, 1% elastane	<1	<5	-
TX12081	Levi's	Hong Kong	Vietnam	jeans	man	100% cotton	<1	<5	-
TX12082	Levi's	Taiwan	Mainland China	t-shirt	man	100% cotton	* 9.7	<5	-
TX12083	Levi's	Thailand	Thailand	denim shirt	woman	100% cotton	<1	<5	-
TX12084	Levi's	Philippines	Mainland China	jeans	man	100% cotton	600	<5	-
TX12085	Levi's	Mexico	Mexico	jeans	woman	99% cotton, 1% elastane	3 100	<5	-
TX12086	Levi's	Switzerland	Turkey	t-shirt	man	100% Cotton	* <1	<5	12
TX12087	Levi's	USA	Mexico	jeans	man	100% cotton	4 100	<5	-
TX12088	Levi's	South Africa	Vietnam	jeans	woman	100% Cotton	5.7	<5	-
TX12089	Levi's	Belgium	Mainland China	t-shirt	man	100% cotton	80	<5	-
TX12090	Levi's	Turkey	Unknown	hoodie	man	unknown	18	<5	-
TX12091	Levi's	Indonesia	Mainland China	t-shirt	man	100% cotton	* <1	<5	138
TX12092	Mango	Philippines	Bangladesh	t-shirt	woman	100% cotton	<1	<5	-
TX12093	Mango	Netherlands	Vietnam	jacket	woman	98% cotton, 2% elastane. lining: 100% polyester	<1	<5	-
TX12094	Mango	Finland	Mainland China	t-shirt	woman	unknown	<1	<5	-
TX12095	Mango	UK	Mainland China	jeans	woman	100% cotton	1 400	<5	-
TX12096	Mango	Spain	Turkey	t-shirt	woman	100% cotton	9 800	<5	-
TX12097	Mango	Austria	Morocco	trousers	woman	100% polyester	7.2	<5	-
TX12098	Mango	South Africa	Turkey	t-shirt	woman	100% cotton	* <1	<5	13
TX12099	Mango	Turkey	Bangladesh	rumper suit	woman	55% linen, 45% cotton	1 500	<5	-
TX12100	Mango	Lebanon	Indonesia	coat	woman	100% polyester	15	<5	-
TX12101	Mango	Israel	Mainland China	dress	woman	100% polyester	1.3	<5	-
TX12102	Marks & Spencer	Thailand	Mainland China	underwear	woman	80% silk, 13% polyamide, 7% elastane	2 100	<5	-
TX12103	Marks & Spencer	Philippines	Indonesia	shorts	man	68% cotton, 32% polyamide	620	<5	-
TX12104	Marks & Spencer	UK	India	underwear	woman	95 % cotton, 5% elastane	<1	<5	-
TX12105	Marks & Spencer	UK	Turkey	pyjama top(pt set)	child	100% cotton	* <1	<5	15
TX12106	Marks & Spencer	Turkey	Turkey	t-shirt	woman	100% linen	84	<5	-
TX12107	Marks & Spencer	Indonesia	Turkey	top	woman	100% polyester	550	<5	-
TX12108	Tommy Hilfiger	Mexico	Mexico	jeans	man	100% cotton	500	<5	-
TX12109	Tommy Hilfiger	Sweden	Vietnam	polo shirt	---	unknown	<1	<5	-
TX12110	Tommy Hilfiger	USA	Philippines	t-shirt	man	100% cotton exclusive of decoration	* 26	<5	200 013

Sample code	Brand	Place of sale	Place of manufacture	Kind of product	Man Woman Child	Fabric	NPEs (mg/kg)	Amines (mg/kg)	Phthalates, total (mg/kg)
TX12111	Tommy Hilfiger	Spain	Turkey	top	woman	80% polyester, 20% viscose excluding decoration	30	<5	-
TX12112	Tommy Hilfiger	Russia	Turkey	jeans	man	100% cotton	17	<5	-
TX12113	Tommy Hilfiger	Russia	Mainland China	shirt	man	100% cotton	<1	<5	-
TX12114	Tommy Hilfiger	Italy	Sri Lanka	t-shirt	woman	96% cotton, 4% elastane excluding decoration	3.9	<5	-
TX12115	Tommy Hilfiger	Austria	Bangladesh	t-shirt	man	100% cotton	* 8.6	<5	376 079
TX12116	Tommy Hilfiger	Canada	Bangladesh	shorts	man	100% cotton exclusive of decoration	<1	<5	-
TX12117	Victoria's Secret	Netherlands	Mainland China	bra	woman	65% nylon, 35% elastane	<1	<5	-
TX12118	Victoria's Secret	Netherlands	Mainland China	bra	woman	satin- 90% nylon, 10 % elastane; embroidery - 62% polyester, 38% nylon excluding decorations	<1	<5	-
TX12119	Victoria's Secret	USA	Sri Lanka	underwear	woman	95% cotton, 5% elastane	* 7.0	<5	5217
TX12120	Victoria's Secret	Canada	Mainland China	camisole top	woman	100% nylon excluding decoration	10	<5	-
TX12121	Zara	Mainland China	Mainland China	jacket	child	100% polyester	2 600	<5	-
TX12122	Zara	Taiwan	Bangladesh	trousers	child	polyurethane fiber	79	<5	-
TX12123	Zara	Germany	India	dress	woman	100% polyurethane	9.6	<5	-
TX12124	Zara	Thailand	Bangladesh	jeans	woman	98% cotton, 2% elastane	<1	<5	-
TX12125	Zara	Denmark	Turkey	t-shirt	---	unknown	* <1	-	4
TX12126	Zara	Spain	Vietnam	coat	child	outershell: 100% polyester. body lining: 65% polyester, 35% cotton. filling: 100% polyester	25	<5	-
TX12127	Zara	Turkey	Spain	shorts	woman	outer-100% cotton; inner-67% polyester, 33% cotton	<1	<5	-
TX12128	Zara	Lebanon	Pakistan	jeans	child	100% cotton	19	<i>o</i> -dianisidine (7 mg/kg)	-
TX12129	Zara	Israel	Morocco	dress	woman	100% polyester	<1	<5	-
TX12130	Zara	Hungary	Pakistan	jeans	child	100% cotton	29	<i>o</i> -dianisidine (9 mg/kg)	-
TX12131	Metersbonwe	Mainland China	Mainland China	skirt	woman	100% viscose rayon	<1	<5	-
TX12132	Metersbonwe	Mainland China	Mainland China	t-shirt	man	100% cotton	* 140	<5	10
TX12133	Metersbonwe	Mainland China	Mainland China	jeans	man	79.1% cotton, 18.7% polyester, 2.2% other	2 100	<5	-
TX12134	Metersbonwe	Mainland China	Mainland China	sweater	man	100% cotton	1 500	<5	-
TX12135	Vancl	Mainland China	unknown	underwear	man	100% cotton	7.6	<5	-
TX12136	Vancl	Mainland China	unknown	t-shirt	woman	50% cotton, 50% modal (modified cellulose)	* 8.5	<5	87
TX12137	Vancl	Mainland China	unknown	cardigan	child	67% polyester, 33% cotton	140	<5	-
TX12138	Vancl	Mainland China	unknown	jeans	man	100% cotton	150	<5	-
TX12139	Blažek	Czech Republic	unknown	jean shorts	man	97% cotton 3% elasten	330	<5	-
TX12140	Blažek	Czech Republic	unknown	shirt	man	75% cotton, 20% PA, 5% EA	47	<5	-
TX12141	Blažek	Czech Republic	unknown	t-shirt	man	50% cotton, 45% modal, 5% elastane	<1	<5	-
TX12142	Blažek	Czech Republic	unknown	underwear	man	50% cotton/45% modal, 5% elastene	<1	<5	-

Table A2: Concentrations of individual phthalates in the 31 articles tested.

Concentrations (mg/kg), in plastisol printed fabric, of the following phthalates; dimethyl phthalate (DMP), diethyl phthalate (DEP), di-n-butyl phthalate (DnBP), diisobutyl phthalate (DIBP), butyl benzyl phthalate (BBP), di-2-ethylhexyl phthalate (DEHP), di-n-octyl phthalate (DnOP), diisononyl phthalate (DINP) and diisodecyl phthalate (DIDP).

Sample code	Brand	Place of sale	Place of manufacture	DMP	DEP	DnBP	BBP	DEHP	DnOP	DINP	DIDP	DIBP
TX12008	Armani	Italy	Turkey	< 3.0	13	17	23 000	200 000	< 3.0	<3.0	<3.0	410
TX12010	Benetton	Mexico	Romania	< 3.0	29	11	55	9.6	< 3.0	< 3.0	< 3.0	23
TX12011	Benetton	Czech Republic	Tunisia	< 3.0	9.4	5.2	7	4.8	< 3.0	< 3.0	< 3.0	6.3
TX12013	Benetton	UK	Bangladesh	< 3.0	11	20	9.1	< 3.0	< 3.0	< 3.0	< 3.0	7.3
TX12025	Only	Mainland China	unknown	< 3.0	3.3	4	< 3.0	5.9	< 3.0	< 3.0	< 3.0	4.8
TX12028	Jack & Jones	Mainland China	unknown	< 3.0	< 3.0	3.7	< 3.0	5.8	< 3.0	< 3.0	< 3.0	4.4
TX12032	Jack & Jones	Lebanon	Bangladesh	< 3.0	3.7	9.8	< 3.0	< 3.0	< 3.0	< 3.0	< 3.0	3.5
TX12040	Calvin Klein	Canada	India	< 3.0	< 3.0	4.7	< 3.0	< 3.0	< 3.0	< 3.0	< 3.0	4.1
TX12041	C&A	Mexico	Mexico	< 3.0	< 3.0	4.5	< 3.0	42	< 3.0	14	< 3.0	< 3.0
TX12044	C&A	Belgium	Unknown	< 3.0	< 3.0	4	< 3.0	13	< 3.0	< 3.0	< 3.0	16
TX12045	C&A	Hungry	Unknown	< 3.0	< 3.0	4.5	8.9	< 3.0	< 3.0	< 3.0	< 3.0	4.4
TX12050	Diesel	Russia	Mainland China	< 3.0	8.5	15	< 3.0	24	< 3.0	< 3.0	< 3.0	8.5
TX12054	Diesel	Israel	Mainland China	< 3.0	8.1	22	< 3.0	16	< 3.0	< 3.0	< 3.0	37
TX12055	Diesel	Hungry	Mainland China	< 3.0	< 3.0	< 3.0	< 3.0	53	< 3.0	< 3.0	< 3.0	3.9
TX12058	Esprit	Germany	Unknown	< 3.0	< 3.0	3.4	< 3.0	5.6	< 3.0	< 3.0	< 3.0	5.3
TX12068	Gap	USA	Indonesia	< 3.0	5.8	4.2	< 3.0	< 3.0	< 3.0	< 3.0	< 3.0	4.4
TX12070	GAP	South Africa	Vietnam	< 3.0	< 3.0	4.5	< 3.0	6.3	< 3.0	< 3.0	< 3.0	3
TX12072	Gap	France	Vietnam	< 3.0	5.8	13	< 3.0	< 3.0	< 3.0	< 3.0	< 3.0	6.5
TX12074	Gap	Canada	Indonesia	< 3.0	18	3.2	< 3.0	< 3.0	< 3.0	< 3.0	< 3.0	4.5
TX12075	H&M	Denmark	Turkey	< 3.0	16	< 3.0	< 3.0	3.8	< 3.0	< 3.0	< 3.0	3
TX12082	Levi's	Taiwan	Mainland China	< 3.0	23	6.5	< 3.0	4.4	< 3.0	< 3.0	< 3.0	< 3.0
TX12086	Levi's	Switzerland	Turkey	< 3.0	3.3	< 3.0	< 3.0	5.1	< 3.0	< 3.0	< 3.0	3.5
TX12091	Levi's	Indonesia	Mainland China	< 3.0	5.8	4	< 3.0	120	< 3.0	< 3.0	< 3.0	7.9
TX12098	Mango	South Africa	Turkey	< 3.0	< 3.0	9.8	< 3.0	< 3.0	< 3.0	< 3.0	< 3.0	3.4
TX12105	Marks & Spencer	UK	Turkey	< 3.0	< 3.0	< 3.0	< 3.0	15	< 3.0	< 3.0	< 3.0	< 3.0
TX12110	Tommy Hilfiger	USA	Philippines	< 3.0	3.6	4.7	< 3.0	< 3.0	< 3.0	200 000	< 3.0	4.6
TX12115	Tommy Hilfiger	Austria	Bangladesh	< 3.0	4.9	21	23	56 000	11	320 000	< 3.0	19
TX12119	Victoria's Secret	USA	Sri Lanka	< 3.0	6.2	3.1	4	5 200	< 3.0	< 3.0	< 3.0	3.4
TX12125	Zara	Denmark	Turkey	< 3.0	< 3.0	< 3.0	< 3.0	3.8	< 3.0	< 3.0	< 3.0	< 3.0
TX12132	Metersbonwe	Mainland China	Mainland China	< 3.0	< 3.0	< 3.0	< 3.0	3.4	< 3.0	< 3.0	< 3.0	6.1
TX12136	VancI	Mainland China	unknown	< 3.0	3	22	< 3.0	58	< 3.0	< 3.0	< 3.0	3.9

Table A3: Additional substances identified using qualitative chemical screening.

Additional substances identified in individual items by qualitative chemical screening, including the total number of compounds isolated and reliably identified for each sample.

Sample code	Brand	No. of chemicals isolated	No. of chemicals reliably identified	Linear alkanes	Benzophenone	Benzyl benzoate	1,1'-Biphenyl	2,6-Di-tert-butyl-4-methyl-phenol	Ethanol, 2-(2-butoxy-ethoxy)- & derivatives*	Nonylphenol	Hexadecanoic acid (& esters*)	Octadec -anoic, -enoic and -adienoic acid (& esters*)	Amyrin	α-Amyrenone	Sitosterol	Cholesterol	Squalene	Others
TX12001	Armani	12	8	6		Y											Y	
TX12002	Armani	13	5	4		Y												
TX12003	Armani	26	13	12		Y												
TX12004	Armani	17	9	9														
TX12005	Armani	26	5	5														
TX12006	Armani	19	7	7														
TX12007	Armani	6	5	3	Y	Y												
TX12008	Armani	9	7	7														
TX12009	Armani	13	1	0								Y						
TX12010	Benetton	15	10	10														
TX12011	Benetton	33	14	11		Y								Y	Y			
TX12012	Benetton	18	12	10		Y									Y			
TX12013	Benetton	11	6	5											Y			
TX12014	Benetton	19	11	9							butyl*				Y			
TX12015	Benetton	21	13	11		Y	Y											
TX12016	Benetton	17	10	8										Y	Y			
TX12017	Benetton	22	10	7							methyl*	methyl*					Y	
TX12018	Benetton	7	5	4										Y				
TX12019	Vero Moda	7	7	7														
TX12020	Vero Moda	0	0	0														
TX12021	Vero Moda	15	11	11														
TX12022	Vero Moda	43	14	13											Y			
TX12023	Vero Moda	9	8	8														
TX12024	Only	19	12	11										Y				
TX12025	Only	1	0	0														
TX12026	Only	12	9	8											Y			
TX12027	Only	16	12	10								methyl*						
TX12028	Jack & Jones	56	15	13										Y			Y	
TX12029	Jack & Jones	20	11	8							methyl*	methyl*		Y				
TX12030	Jack & Jones	28	14	12										Y			Y	
TX12031	Jack & Jones	26	11	11														
TX12032	Jack & Jones	17	12	11										Y				
TX12033	Calvin Klein	20	6	6														
TX12034	Calvin Klein	20	15	13		Y					octadecyl*							
TX12035	Calvin Klein	46	14	13										Y				
TX12056	Esprit	11	1	0														
TX12057	Esprit	7	7	7														octyl-diphenylamine
TX12058	Esprit	19	6	6														
TX12059	Esprit	13	7	7														

Sample code	Brand	No. of chemicals isolated	No. of chemicals reliably identified	Linear alkanes	Benzophenone	Benzyl benzoate	1,1'-Biphenyl	2,6-Di-tert-butyl-4-methyl-phenol	Ethanol, 2-(2-butoxy-ethoxy)- & derivatives*	Nonylphenol	Hexadecanoic acid (& esters*)	Octadec -anoic, -enoic and -adienoic acid (& esters*)	Amyrin	α-Amyrenone	Sitosterol	Cholesterol	Squalene	Others
TX12060	Esprit	27	9	7		Y									Y			
TX12061	Esprit	17	8	6										Y	Y			
TX12062	Esprit	14	3	2											Y			
TX12063	Esprit	35	14	11		Y									Y			benzenemethanamine, N-(phenylmethyl-
TX12064	Esprit	14	5	4								methyl*						
TX12081	Levi's	24	11	8									Y	Y	Y			
TX12082	Levi's	15	9	7						Y					Y			
TX12083	Levi's	61	14	11		Y								Y				Ethanol, 2-(tetradecyloxy)-
TX12084	Levi's	38	17	11		Y							Y	Y	Y	Y	Y	
TX12085	Levi's	32	11	9							isopropyl*							Bis(2-ethylhexyl) maleate
TX12086	Levi's	24	10	7										Y	Y	Y		
TX12087	Levi's	38	14	10	Y							isopropyl*		Y	Y			
TX12088	Levi's	31	13	9					Y					Y	Y	Y		
TX12089	Levi's	32	10	9											Y			
TX12090	Levi's	15	10	10														
TX12091	Levi's	22	5	5			Y											
TX12121	Zara	56	15	13														Benzene, 1,1'-(3-methyl-1-propene-1,3-diyl)bis-
TX12122	Zara	58	13	9				Y						Y	Y			Benzyl naphthyl ether
TX12123	Zara	18	5	3							methyl*	methyl*						
TX12124	Zara	60	11	11														
TX12125	Zara	44	13	11										Y	Y			
TX12126	Zara	30	9	8				Y										
TX12127	Zara	35	13	7					acetate*	Y	Y	Y		Y	Y			
TX12128	Zara	28	4	3										Y				